

# Physiology of the Hypothalamic-Pituitary Thyroidal System

Last Updated: June 1, 2011

## Authors

- **Stefano Mariotti, M.D.**

Department of Endocrinology Institute of Internal Medicine University of Cagliari Policlinico Universitario S.S. 554, Bivio per Sestu I-09042 Monserrato CA, ITALY Tel: 011-39-70-602-86430 Fax: 011-39-70-602-86429

The activity of the thyroid gland is predominantly regulated by the concentration of the pituitary glycoprotein hormone, thyroid stimulating hormone (TSH). In the absence of pituitary or of thyrotroph function hypothyroidism ensues. Thus, regulation of thyroid function in normal individuals is to a large extent determined by the factors which regulate the synthesis and secretion of TSH. Those factors are reviewed in this chapter and consist principally of thyrotropin releasing hormone (TRH) and the feedback effects of circulating thyroid hormones at the hypothalamic and pituitary levels. The consequence of the dynamic interplay of these two dominant influences on TSH secretion, the positive effect of TRH on the one hand and the negative effects of thyroid hormones on the other, result in a remarkably stable concentration of TSH in the circulation and consequently little alteration in the level of circulating thyroid hormones from day to day and year to year. This regulation is so carefully maintained that an abnormal serum TSH in most patients is believed to indicate the presence of a disorder of thyroid gland function. The utility of TSH measurements has been recognized and its use has remarkably increased, due to the development of immunometric methodologies for its accurate quantitation in serum, although the criteria to define a "normal range" still remain matted of controversy.

This chapter is organized into two general sections. The first portion reviews basic studies of TSH synthesis, post-translational modification and release. The second deals with physiological studies in humans which serve as the background to the diagnostic use of TSH measurements and reviews the results of TSH assays in a pathophysiological context.

## **The Regulation of Thyroid-Stimulating-Hormone synthesis and secretion: Molecular Biology and Biochemistry**

### **The TSH Molecule**

TSH is a heterodimer consisting of  $\alpha$  and  $\beta$  subunits tightly, but non-covalently, bound.<sup>(38)</sup> While the molecular weight of the deduced amino acid sequence of mature  $\alpha$  plus TSH  $\beta$  subunits is approximately 28,000 Da, additional carbohydrate (15% by weight) results in a significantly higher molecular weight estimate based on sizing by polyacrylamide gel electrophoresis. The  $\alpha$  subunit is

common to TSH, follicle stimulating hormone (FSH), luteinizing hormone (LH), and chorionic gonadotropin (CG). The  $\beta$  subunit confers specificity to the molecule since it interacts with the thyroid cell TSH receptor (TSH-R) and is rate-limiting in the formation of the mature heterodimeric protein. However, the free  $\beta$  subunit is inactive and requires noncovalent combination with the  $\alpha$  subunit to express hormonal bioactivity. The linear sequence of human  $\alpha$  subunit is represented by 92 amino acids including 10 half-cystine residues, all of which in disulfide linkage. The human TSH  $\beta$  (hTSH  $\beta$ ) subunit contains 118 amino acids, as predicted by complementary DNA sequences, but hTSH  $\beta$  isolated from pituitary gland has an apoprotein core of 112 amino acids, due to carboxyl-terminal truncation during purification.

The production rate (PR) of human TSH is normally between 50 and 200 mU/day and increases markedly (up to >4000 mU/day) in primary hypothyroidism; the metabolic clearance rate (MCR) of the hormone is about 25 ml/min/m<sup>2</sup> in euthyroidism, while is significantly higher in hypothyroidism and lower in hyperthyroidism.<sup>(3)</sup> The PR of free  $\alpha$  subunit is about 100  $\mu$ g/day, increases only about twice in primary hypothyroidism and/or in post-menopausal women and decreases (about to one half) in hyperthyroidism.<sup>(4)</sup> The PR of free TSH  $\beta$  subunit is too low to be calculated in all hyperthyroid and in most euthyroid subjects, while is 25-30  $\mu$ g/day in primary hypothyroidism.<sup>(4)</sup> The MCR of free subunits is 2-3 times faster than that of TSH, being about 68 ml/min/m<sup>2</sup> for  $\alpha$  and 48 ml/min/m<sup>2</sup> for  $\beta$  subunit.<sup>(4)</sup>

The human  $\alpha$  subunit gene is located on chromosome 6 and the TSH  $\beta$  gene on chromosome 1.<sup>(5)</sup> The structure of  $\alpha$  subunit gene has been determined in several animal species.<sup>(38)</sup> The genes of each species are approximately of the same size and similarly organized in four exons and three introns. The human gene is 9.4 kilobases (kb) in length, with three introns measuring 6.4 kb, 1.7 kb and 0.4 kb, respectively. The TSH  $\beta$  subunit gene has been isolated in mouse<sup>(6)</sup>, rat<sup>(8)</sup>, and humans.<sup>(38)</sup> At difference with the  $\alpha$  subunit, the organization of the TSH  $\beta$  gene is somewhat variable between the different species. The rat and the human genes are organized in three exons, while the mouse gene contains two additional 5'-untranslated exons. The first exon is untranslated, the leader peptide and the first 34 aminoacids are encoded by the second exon, while the third exon represents the remaining coding region and 3'-untranslated sequences. A single transcription start has been identified in hTSH  $\beta$  gene, while the rat and the mouse genes contain two starting sites separated by approximately 40 base pairs (bp); most transcription begins from the downstream site, which corresponds to the location of the human transcriptional start. A schematic representation of TSH  $\beta$  gene is reported in Fig. 4.1.




Figure 4-1 Thyrotropin  $\beta$  (TSH  $\beta$ ) gene structure and mutations found in patients with congenital central hypothyroidism (modified from McDermott et al. (36) and Baquedano et al (38))

The pre-translational regulation of TSH synthesis and secretion is a complex process, detailed in the next paragraphs. The formation of mature TSH involves several post-translational steps including the

excision of signal peptides from both subunits and co-translational glycosylation with high mannose oligosaccharides.<sup>(38)</sup> As the glycoproteins are successively transferred from the rough endoplasmic reticulum to the golgi apparatus, the trimming of mannose and further addition of fucose, galactose and sialic acid occurs.<sup>(13)</sup> The  $\alpha$  subunit has two and TSH  $\beta$  one asparagine (N)-linked oligosaccharides showing typical biantennary structure fully sulfated in bovine and half-sulfated in human TSH.<sup>(2)</sup> The primary intracellular role of these glycosylation events may be to allow proper folding of the  $\alpha$  and TSH  $\beta$  subunits permitting their heterodimerization and also preventing intracellular degradation.<sup>(38)</sup> On the basis of crystallographic studies on hCG and other glycoprotein hormones, an homology model of the tridimensional structure of TSH has been proposed.<sup>(15)</sup> This model (Fig. 4-2) predicts for both  $\alpha$  and  $\beta$  subunits the presence of two  $\beta$ -hairpin loops (L1 and L3) on one side of a central “cystine (pair of cysteine molecules) knot” formed by three disulfide bonds, and a long loop (L2) on the other side. Both  $\alpha$  and  $\beta$  chains have functionally important domains involved in TSH-R binding and activation (Fig. 4-2). Of particular relevance is the so-called “seat belt” region of the  $\beta$  chain comprised between the 10<sup>th</sup> (C86) and the 12<sup>th</sup> (C105) cystein residue (Fig. 4-2 and Fig. 4-3). The name “seat belt” derives from the conformational structure of the  $\beta$  chain determined by the disulfide bridge (C39/C125) toward the C-terminal tail of the  $\beta$  subunit that wraps the  $\alpha$  subunit like a “setas belt” (Fig. 4.3) and stabilizes the heterodimerization of TSH.<sup>(38)</sup> Proper TSH glycosylation is also necessary to attain normal bioactivity,<sup>(17)</sup> a process which requires the interaction of the neuropeptide thyrotropin releasing hormone (TRH, Fig. 4-4), with its receptor on the thyrotroph.<sup>(38)</sup> The requirement for TRH in this process is illustrated by the fact that in patients with central hypothyroidism due to hypothalamic-pituitary dysfunction, normal or even slightly elevated levels of radioimmunoassayable, but biologically subpotent TSH are found in the circulation in the presence of a reduced free T4.<sup>(38)</sup> Chronic TRH administration to such patients normalized the glycosylation process enhancing both its TSH-R binding affinity as well as its capacity to activate adenyl cyclase. This, in turn, can normalize thyroid function in such patients.<sup>(23)</sup> On the other hand, enhanced TSH bioactivity is invariably found in sera from patients with thyroid hormone resistance.<sup>(24)</sup> Moreover, variations of TSH bioactivity (mostly related to different TSH glycosylation) have been observed in normal subjects during the nocturnal TSH surge, in normal fetuses during the last trimester of pregnancy, in primary hypothyroidism, in patients with TSH-secreting pituitary adenoma and in non-thyroidal illnesses.<sup>(38)</sup> Glycosylation of the molecule can also influence the rapidity of clearance of TSH from the circulation. Taken together, these findings have lead to a new concept of a qualitative regulation of TSH secretion, mainly achieved through both the transcriptional and posttranscriptional mechanisms involved in TSH glycosylation.<sup>(26)</sup>




Figure 4-2 Schematic drawing of human TSH, based on a molecular homology model built on the template of a hCG model 14 . The  $\alpha$ -subunit is shown as checkered, and the  $\beta$ -subunit as a solid line. The two hairpin loops in each subunit are marked L1, L3; each subunit has also a long loop (L2), which extends from the opposite site of the central cystine knot. The functionally important  $\alpha$ -subunit domains are boxed. Important domains of the  $\beta$ -subunit are marked directly within the line drawing (crossed line, beaded line and dashed line): For further details the reader is referred to Grossman et al. 2 . (Reproduced from Grossman et al. (2) , with permission).




Figure 4-3 (new) Structural model of TSH based on the FSH x-ray structure, which is the best available structural template for TSH. The boxed residue numbers represent cysteines residues, which form stabilizing disulfide bridges (yellow): 5 in alpha-subunit (red orange), and 6 in the beta-subunit (magenta). The disulfide bridge (C39/C125) toward the C-terminal tail of the beta-subunit of TSH that wraps around the beta-subunit like a “seat belt” stabilizes the heterodimerization of TSH as well and also in FSH, LH, and CG. (Reproduced from Kleinau et al.(16), with permission)




Figure 4-4 Structure of TRH.

Specific amino acid sequences in the common  $\alpha$  and TSH  $\beta$  subunits are critical for the heterodimerization, secretion and bioactivity of mature TSH. These sequences include highly conserved segments which are essential for TSH-R binding and biological activity (see Refs <sup>(38)</sup> for an extensive review). The peptide sequence <sup>27</sup>CAGYC<sup>31</sup> (cysteine-alanine-glycine-tyrosine-cysteine) is highly conserved in TSH  $\beta$ , LH  $\beta$ , hCG  $\beta$  and FSH  $\beta$  and is thought to be important in subunit

combination .<sup>(38)</sup> Several inherited TSH  $\beta$  gene mutations responsible for familial isolated central hypothyroidism are listed in Table 4-1 and depicted in Fig. 4-1. The most frequent mutation is a homozigous single base deletion in codon 105 (C105D, 114X) leading to unstable heterodimer.<sup>(38)</sup>

Table 4-1 Mutations of the TSH  $\beta$  gene responsible of isolated central hypothyroidism: effects on TSH heterodimer formation.

Mutation of TSH $\beta$ gene	Consequence of mutation on TSH heterodimer formation
G29R <sup>(38)</sup>	Prevents dimer formation modifying the CAGYC region
E12X <sup>(29)</sup>	Truncated TSH $\beta$ subunit unable to associate with $\alpha$ chain
C105V, 114X <sup>(38)</sup>	Change of amino acid sequence in the “seat belt” region leading to unstable heterodimer
Q49X <sup>(38)</sup>	Truncated TSH $\beta$ subunit forming a bio-inactive heterodimer with the $\alpha$ chain
IVS2+5 à A <sup>(35)</sup>	Base substitution at intron 2 (position +5) with shift of the translational start point to an out of frame position of exon 3 resulting in a truncated transcript
C85R <sup>(39)</sup>	T to C transition at codon 85 of exon 3, resulting in a change of cysteine to arginine, preventing the formation of a functional heterodimer with the $\alpha$ -subunit
C162G à A <sup>(38)</sup>	G to A change at the 5' donor splice site of exon/intron 2 transition causing a (CGA à CGG) polymorphisms, which although <i>per se</i> silent, disrupts the 5' consensus sequence critical for slicing causing complete exon 2 skipping
C88Y <sup>(38)</sup>	The loss of one of the cysteine residues critical for $\beta$ -subunit conformation and intracellular degradation

The understanding of the relationship between molecular structure and biological activity of TSH recently allowed the synthesis of TSH variants designed by site-directed mutagenesis with either antagonist<sup>(39)</sup> or superagonist<sup>(38)</sup> activity that potentially offer novel therapeutic alternatives .

## Other thyrotropic hormones

Recently, a new thyrotropic hormone represented by a heterodimer of two new glycoprotein subunits (glycoprotein hormone  $\alpha$ 2-subunit – GPA2 and glycoprotein hormone  $\beta$ 5-subunit – GPB5) has been identified in human pituitary and called thyrostimulin.<sup>(41)</sup> Thyrostimulin has a sequence similarity of 29% with  $\alpha$ - and 43% with  $\beta$ -subunit and is able to activate TSH-R.<sup>(38)</sup> Although it has been hypothesized that it could account for the residual stimulation of thyroid gland observed in patients with central hypothyroidism,<sup>(43)</sup> its physiological role is still unknown. In the GPA2/GPB5 heterodimer is localized in extrapituitary tissues such eye, testis and ovary,<sup>(38)</sup> while anterior pituitary express almost exclusively GPA2.<sup>(38)</sup> In rat ovary thyrostimulin activate TSH-R expressed by granulosa cells,<sup>(38)</sup> indicating a potential paracrine activity.

## Regulation of TSH Synthesis and Secretion.

The major regulators of TSH production are represented by the inhibitory effects of thyroid hormone<sup>(45)</sup> and by the stimulatory action of TRH. As shown in Fig. 4-5, T3 acts via binding to the thyrotropin nuclear T3 receptor, and T4 mainly acts via its intra-pituitary or intra-hypothalamic conversion to T3,

although a direct negative effect of T<sub>4</sub> independent from local T<sub>3</sub> generation on TSH  $\beta$  gene expression.<sup>(38)</sup> Both thyroid hormones regulate directly regulate the synthesis and release of TSH at the pituitary level and indirectly affect TSH synthesis via their effects on TRH and other neuropeptides. TRH is the major positive regulator of hTSH  $\beta$  gene expression and mainly acts by activating the phosphatidylinositol-protein C pathway. Other hormones/factors are also implicated in the complex regulation of TSH- $\beta$  gene expression, as detailed below.




Figure 4-5 Basic elements in the regulation of thyroid function. TRH is a necessary tonic stimulus to TSH synthesis and release. TRH synthesis is regulated directly by thyroid hormones. T<sub>4</sub> is the predominant secretory product of the thyroid gland, with peripheral deiodination of T<sub>4</sub> to T<sub>3</sub> in the liver and kidney supplying roughly 80% of the circulating T<sub>3</sub>. Both circulating T<sub>3</sub> and T<sub>4</sub> directly inhibit TSH synthesis and release independently; T<sub>4</sub> via its rapid conversion to T<sub>3</sub>. SRIH = somatostatin.

## Effects of thyroid hormone on TSH synthesis and release

In animal models, thyroid hormone administration is followed by a marked decrease of both  $\alpha$  and TSH  $\beta$  subunit mRNA,<sup>(38)</sup> but TSH  $\beta$  is suppressed more rapidly and more completely than  $\alpha$  subunit. In humans with primary hypothyroidism a paradoxical increase of serum TSH concentration has been observed shortly after beginning thyroid hormone replacement therapy, followed later by TSH suppression.<sup>(49)</sup> The precise mechanism for this phenomenon has not been fully elucidated: it could be due to a generalized defect in protein synthesis as a consequence of hypothyroidism, or to the presence of a still unrecognized stimulatory thyroid hormone cis-acting element (see below). Thyroid hormone regulation of TSH  $\beta$  subunit transcription is complex and, at least in the rat and mouse, involves control of gene transcription at both start sites of the gene<sup>(38)</sup> (Fig. 4-6). Studies of the human, rat and mouse TSH  $\beta$  genes have demonstrated that they contain DNA hexamer half sites with strong similarity to the T<sub>3</sub> response elements (TREs) found in genes which are positively regulated by thyroid hormone<sup>(38)</sup> (see Chapter 3). The sequences in the TSH  $\beta$  gene are diagrammed in Fig. 4-6 and their similarity to the typical hexamer binding sites in positively regulated genes and in the rat  $\alpha$  subunit gene is seen by comparison to the TRE sequences from positively regulated genes<sup>(60)</sup> (see Chapter 3). In keeping with this concept, T<sub>3</sub> exerts similar negative activity on rat GH3 cells transfected with plasmids constructs containing the putative negative TRE of rat TSH  $\beta$  or containing a half-site motif of the consensus

positive TRE. <sup>(38)</sup> The conserved TRE-like sequences are the best candidate site on the TSH gene to which the T3 receptor (TR) binds. The subsequent binding of T3 to TR-DNA complexes suppresses transcription of both  $\alpha$  and TSH  $\beta$  subunit genes. <sup>(38)</sup> The inhibitory effect of thyroid hormone is observed with all  $\alpha$  and  $\beta$  isoforms of TR, but TR- $\beta$ 2 (a TR isoform with pituitary and central nervous system-restricted expression) has the greatest effect. <sup>(38)</sup> This *in vitro* observation is in keeping with a series of *in vivo* data obtained in transgenic and knockout mice with generalized or pituitary-selective expression of mutated TR isoform genes. Knockout mice for TR- $\alpha$ 1 develop only minor abnormalities in circulating T4 and TSH concentration <sup>(65)</sup>, while mice lacking both  $\beta$ 1 and  $\beta$ 2 isoforms ( $\beta$ -null) develop increased serum T4 and TSH level, but retain partial TSH suppression by T3 administration. <sup>(38)</sup> Mice selectively lacking TR- $\beta$ 2 isoform develop hormonal abnormalities similar to TR- $\beta$ -null animals, indicating a key role of TR- $\beta$ 2 as mediator of T3-dependent negative regulation. <sup>(68)</sup> On the other hand, the residual T3-dependent TSH suppression observed in mice lacking TR- $\beta$  isoforms suggests that TR- $\alpha$ 1 may partially substitute for TR- $\beta$  in mediating T3 suppression: accordingly, mice lacking all ( $\alpha$  and  $\beta$ ) TR isoforms develop dramatic increase in circulating T4 and TSH concentration, indicating that a complete expression of all TR isoforms are required for normal regulation of the hypothalamic-pituitary thyroid axis. <sup>(38)</sup> Further studies have been carried out with models of mice expressing selectively at the pituitary <sup>(38)</sup> and hypothalamic <sup>(73)</sup> level different combinations of double homozygous or combined heterozygous deletions of both TR- $\alpha$  and TR- $\beta$  genes. These studies confirmed the key role of TR- $\beta$  integrity both at the pituitary and hypothalamic level for the inhibition of TSH- $\beta$  and TRH gene expression. TR- $\alpha$  however, may partially substitute for TR- $\beta$  in mediating a partial thyroid hormone dependent TSH suppression.




Figure 4-6 DNA sequences of the putative TREs in the rat, mouse, and human TSH  $\beta$  subunit gene promoters. A comparison of the proximal promoter regions of the rat, mouse, and human TSH  $\beta$  subunit genes is shown. The straight arrows denote TRE consensus half-sites identified by functional and TR binding assays. The first exons (relative to the downstream promoter for the rat and mouse genes) are shaded, and the bent arrows denote the sites of transcription initiation. Note a nine-nucleotide deletion in the human gene relative to the rodent genes indicated by the triangle just 5'

The negative transcription conferred by TSH  $\beta$  TRE sequences is retained even if they are transferred to a different gene or placed in a different position within a heterologous gene. <sup>(38)</sup> This suggests that the negative transcriptional response to thyroid hormone is intrinsic to this TRE structure. In contrast with positive TREs, little is known about the mechanism of T3-dependent negative regulation of genes like TSH  $\beta$ . Data discussed above clearly show the crucial role of the TR- $\beta$  in the negative regulation of TSH synthesis. Like positive TREs, it has been recently established that TR binding to DNA is required for negative gene regulation. <sup>(77)</sup> Early experiments suggested that unliganded TR homodimers stimulate the expression of TSH  $\beta$ , (a behavior appearing a mirror image of the silencing effect on positive TREs), but the methodology employed was not adequate to study the low level of basal TSH  $\beta$  transcriptional activity. More recently, the use of CV1 cell lines containing the TSH  $\beta$ -CAT (chloramphenicol acetyltransferase) reporter allowed a more accurate study of the molecular

mechanisms involved in the liganded TR suppression. (38) In this experimental system TSH  $\beta$  gene suppression was dependent on the amounts of T3 and TR, but unliganded TR did not stimulate TSH  $\beta$  activity, suggesting that TR itself is not an activator. Moreover recruiting of co-activators and co-repressors were shown to be not necessarily essential, but required for full suppression of TSH  $\beta$  gene. (38)

In contrast to the potentiating activity exerted on stimulatory TREs, retinoid X receptors (RXR) either unliganded or in combination with retinoic acid (RA) block thyroid hormone-mediated inhibition of TSH  $\beta$  gene, possibly through competition with the TR-T3 complex binding to DNA. (38) However , RA is also able to suppress TSH  $\beta$  gene production when bound to RAR and RXR interacting with response elements separate from negative TREs. (38) Taken together, these findings imply that distinct mechanisms are involved in thyroid hormone dependent inhibition and stimulation of TSH synthesis. Indirect support to this concept derives from the identification of patients with selective pituitary thyroid hormone resistance carrying TR mutations associated with normal or enhanced function on stimulatory TRE in peripheral tissues, but defective function on inhibitory TREs of TSH  $\beta$  and TRH genes. (83)

An other peculiar feature of the negative TSH  $\beta$  TRE is that its 5' portion (Fig 4-6 and 4-13) displays high homology with the consensus sequence of binding sites for c-Jun and c-Fos, which heterodimerize to form the transcription factor called AP-1. This makes the negative TSH  $\beta$  TRE a "composite element" able to bind both thyroid hormone receptors and AP-1. (38) Since AP-1 antagonizes *in vitro* the inhibition exerted by thyroid hormone, it may act *in vivo* as a modulator of TRH-dependent regulation of TSH  $\beta$  gene .(38) The role of other important TSH  $\beta$  gene activity modulators (such as Pit-1 and its splicing variants) will be discussed later. Other abnormalities of the mechanisms involved in the negative feed-back on TSH by thyroid hormones could be involved in rare pathological conditions of difficult identification and diagnosis.

Since unliganded TR does not behave as an activator of TSH  $\beta$  gene, other mechanisms are involved in the increase of TSH production observed in hypothyroidism. In the hypothyroid rat the TSH production is increased 15 to 20 fold over that in the euthyroid state. This can be attributed to the stimulatory effects of TRH (see below) unopposed by the negative effects of T3; moreover, besides the transcription rate per cell, there is a 3 to 4 fold increase in the absolute number of thyrotrophs in the hypothyroid pituitary. (86) Electron microscopic studies have shown near total depletion of secretory granules in the thyrotrophs of hypothyroid animals, a change that is reversed soon after administration of thyroid hormone. (87)

## Thyroid hormone effects on release of TSH

The acute administration of T3 to the hypothyroid rat causes a rapid and marked decrease in the level of serum TSH. (38) (Fig. 4-7) This decrease occurs prior to the decrease in pituitary  $\alpha$  and TSH  $\beta$  mRNAs. (38) . During the period that circulating TSH is falling, pituitary TSH content remains unchanged or increases slightly. (91) The suppression of TSH release is rapid, beginning within 15 minutes of intravenous T3 injection, but is preceded by the appearance of T3 in pituitary nuclei. (91) In the experimental setting in the rat, as the bolus of injected T3 is cleared and the plasma T3 level falls, nuclear T3 decreases followed shortly by a rapid increase in plasma TSH. Both the chronological and quantitative relationships between receptor bound T3 and TSH release are preserved over this time. (91)




Figure 4-7 Time course of specific pituitary nuclear T<sub>3</sub> binding and changes in plasma TSH in hypothyroid rats after a single intravenous injection of 70 ng T<sub>3</sub> per 100 g of body weight. Since the maximal capacity of thyroid hormone binding in pituitary nuclear proteins is about 1 ng T<sub>3</sub>/mg DNA, the peak nuclear T<sub>3</sub> content of 0.44 ng T<sub>3</sub>/mg corresponds to 44% saturation. The plasma level falls to about 55% of its initial basal level by 90 minutes after T<sub>3</sub> injection demonstrating that there is both a chronological and a quantitative correlation between nuclear T<sub>3</sub> receptor saturation and suppression of TSH release. (From Silva and Larsen, (384) with permission)

The mechanism for this effect of T<sub>3</sub> is unknown. As discussed before, Suppression of basal TSH release is difficult to study *in vitro*. Accordingly, the T<sub>3</sub> induced blockade of TRH-induced TSH release has been used as a model for this event. This T<sub>3</sub> effect is inhibited by blockers of either protein or mRNA synthesis.<sup>(38)</sup> The effect is not specific for TRH since T<sub>3</sub> will also block calcium ionophore, phorbol ester or potassium-induced TSH release.<sup>(38)</sup> Furthermore, T<sub>3</sub> will also block the TRH-induced increase in intracellular calcium which precedes TSH release.<sup>(96)</sup> Thus, T<sub>3</sub> inhibits TSH secretion regardless of what agent is used to initiate that process.

T<sub>4</sub> can cause an equally rapid suppression of TSH via its intrapituitary conversion to T<sub>3</sub>.<sup>(88)</sup> (Fig. 4-5) This T<sub>4</sub> to T<sub>3</sub> conversion process is catalyzed by the Type 2 deiodinase. (see Chapter 3). An effect of T<sub>4</sub> *per se* can be demonstrated if its conversion to T<sub>3</sub> is blocked by a general deiodinase inhibitor such

as iopanoic acid.<sup>(38)</sup> In this case, the T4 in the cell rises to concentrations sufficient to occupy a significant number of receptor sites even though its intrinsic binding affinity for the receptor is only 1/10 that of T3. A similar effect can be achieved by rapid displacement of T4 from its binding proteins by flavonoids.<sup>(98)</sup> It seems likely, however, that under physiological circumstances the feedback effects of T4 on TSH secretion and synthesis can be accounted for by its intracellular conversion to T3.

The effect of suppressive doses of T3, T4 and triiodoacetic acid on serum TSH has been evaluated in humans by ultrasensitive TSH assays.<sup>(99)</sup> TSH suppression was shown to be a complex, biphasic, nonlinear process, with three temporally distinct phases: phase 1, a rapid TSH suppression, starting after 1 h and lasting for 10-20 h; phase 2, slower suppression, starting between 10 and 20 h and lasting for 6-8 weeks; and phase 3, with stable low TSH level (<0.01 mU/L). This pattern of thyroid hormone suppression of TSH is reproducible and independent of the basal thyroid status or the thyroid hormone analog used.

Based on the analyses of the sources of nuclear T3 in the rat pituitary (see Chapter 3), one would predict that approximately half of the feedback suppression of TSH release in the euthyroid state can be attributed to the T3 derived directly from plasma; the remainder accounted for by the nuclear receptor bound T3 derived from intrapituitary T4 to T3 conversion.<sup>(88)</sup> Various physiological studies in both rats and humans confirm this concept in that a decrease in either T4 or T3 leads to an increase in TSH. The effect of T4 is best illustrated in the iodine deficient rat model (Fig. 4-8). In this paradigm, rats are placed on a low iodine diet and serum T3, T4, and TSH quantitated at frequent times thereafter.<sup>(100)</sup> Despite the fact that serum T3 concentrations remain constant, there is a marked increase in TSH as the serum T4 falls. In humans, severe iodine deficiency produces similar effects.<sup>(101)</sup> The most familiar example of the independent role of circulating T4 in suppression of TSH is found in patients in the early phases of primary hypothyroidism in whom serum T4 is slightly reduced, serum T3 is normal or even high normal range, but serum TSH is elevated<sup>(38)</sup> (Table 4-2).




Figure 4-8 Serum T<sub>3</sub>, T<sub>4</sub>, and TSH concentrations (mean  $\pm$  SD) in rats receiving a low iodine diet (LID), with or without potassium iodide (KI) supplementation in the drinking water. (From Riesco et al, (100) with permission)

Table 4-2 Serum concentration of thyroid hormones and TSH in patients with primary hypothyroidism of increasing severity.

			TSH (mU/L)	
	T4	T3	Basal	After 200 µg TRH
Group <sup>a</sup>	( µ g/dl)	(ng/dl)		
Control	7.1 ± 0.9	115 ± 31	1.3 ± 0.5	11 ± 4.6
1	6-9	119 ± 40	5.3 ± 2.3	39 ± 15
2	4-6	103 ± 20	13 ± 10	92 ± 50
3	2-4	101 ± 35	63 ± 56	196 ± 120
4	<2	43 ± 28	149 ± 144	343 ± 326

Results are mean ± SD.

<sup>a</sup> Patients were categorized according to the severity of thyroid disease based on serum T4 concentrations.

(Adapted from Bigos et al, <sup>(103)</sup> with permission)

### The Role of Thyrotropin Releasing Hormone (TRH) in TSH Secretion.

TRH is critical for the synthesis and secretion of TSH either in the presence or absence of thyroid hormones. Destruction of the parvo-cellular region of the rat hypothalamus, which synthesizes the TRH relevant for TSH regulation, causes hypothyroidism. <sup>(38)</sup> Hypothalamic TRH synthesis is in turn regulated by thyroid hormones and thus TRH synthesis and release is an integral part of the feedback loop regulating thyroid status (see Fig. 4-5). TRH also interacts with thyroid hormone at the thyrotroph raising the set-point for thyroid hormone inhibition of TSH release. <sup>(104)</sup> The data supporting these general concepts are reviewed in subsequent sections.

### Control of thyrotroph-specific TRH synthesis

TRH is synthesized as a large pre-pro-TRH protein in the hypothalamus and in several tissues, such as the brain, the β cells of the pancreas, the C cells of the thyroid gland, the myocardium, reproductive organs including the prostate and testis, in the spinal cord and in anterior pituitary. <sup>(38)</sup> Recent investigations employing sophisticated techniques such as fast atom bombardment mass spectrometry and gas phase sequence analysis showed that most TRH immunoreactivity found in extrahypothalamic tissues is actually accounted by TRH-immunoreactive peptides displaying different substitutions of the aminoacid histidine of authentic TRH, which could be active in autocrine/paracrine networks involving also extrapituitary TSH secretion. <sup>(110)</sup> On the other hand, the pituitary TSH production is dependent only on TRH synthesized in specific areas of the paraventricular nucleus (PVN) (Fig. 4.9), located at the dorsal limits of the third ventricle. <sup>(38)</sup> In particular, TRH neurons are almost exclusively found in the parvicellular part of PVN and, while TSH-synthesizing neurons are found in all parvicellular subdivisions of PVN, hypophysiotropic TRH neurons are located exclusively in the periventricular and medial subdivisions (Fig 4.9). Hypophysiotropic TRH neurons project their axons to the median eminence, where TRH is released and drained to the anterior pituitary through the long portal veins. <sup>(38)</sup> Although paracrine and autocrine activity has been recently described for TRH secreted in the anterior pituitary, <sup>(113)</sup> the physiological relevance of pituitary TRH is unknown. The human pre-pro-TRH molecule is a protein of 29 kDa containing 5 progenitor sequences for TRH. <sup>(38)</sup> These five peptides consist of a gln-his-pro-gly peptide preceded and followed by lys-arg or arg-arg di-peptides. The basic di-peptides are the cleavage sites for release of the tetra-peptide progenitor sequence. The glycine

residue is the source of the terminal amide for the proline residue of TRH (Fig. 4-4). In addition to the pro-TRH peptides which are released from the pre-pro TRH molecule, intervening non-TRH peptides which have potential physiological function are co-released.<sup>(38)</sup> In particular, the prepro-TRH fragment 160-169, also known as hST10, TRH-enhancing peptide and Ps4<sup>(38)</sup> is able to stimulate TSH  $\beta$  gene expression and to enhance the TRH-induced release of TSH and prolactin (PRL) from the pituitary.<sup>(38)</sup> Ps4 high affinity receptors have been recently shown within several extrapituitary neural tissues and other endocrine systems (mainly in pancreas and in male reproductive system), and targeted pre-pro TRH gene disruption results in hyperglycemia besides the expected hypothyroidism.<sup>(118)</sup> An other pre-proTRH peptide (fragment 178-199)<sup>(38)</sup> appears to be a modulator of ACTH secretion, although the physiological relevance of this phenomenon is unknown. The prepro-TRH processing is mostly mediated by the prohormone convertases PC1 and PC2, and takes place during axonal transport after removal of the signal peptide.<sup>(124)</sup> Subsequent cleavages occur as the peptides move down the axon toward the nerve terminal, from which TRH is released into the hypothalamic-pituitary portal plexus.<sup>(38)</sup>




Figure 4.9 (new) Distribution of TRH-synthesizing neurons in the PVN. Low power micrographs (A–C) illustrate the TRH neurons at three rostrocaudal levels of the PVN. Schematic drawings (D–F) illustrate the subdivisions of the PVN where hypophysiotropic TRH neurons are localized (gray). AP, anterior parvocellular subdivision; DP, dorsal parvocellular subdivision; LP, lateral parvocellular subdivision; MN, magnocellular part of PVN; MP, medial parvocellular subdivision, PV,

periventricular parvocellular subdivision; III, third ventricle. (From Fekete & Lechan, 2007, (112) with permission)

Thyroid hormones exert strong negative regulation of TRH synthesis at hypothalamic level.<sup>(38)</sup> Increases in TRH mRNA levels occur during primary or secondary hypothyroidism and decreases in TRH mRNA result from implantation of a small crystal of T3 adjacent to the PVN<sup>(127)</sup> This regulation is observed *in vivo* exclusively in the parvo-cellular division of the PVN<sup>(38)</sup> (whose neurones contain functional TR isoforms  $\alpha 1$ ,  $\beta 2$  and  $\beta 1$ <sup>(130)</sup>), while in tissues outside the central nervous system expressing the TRH gene, negative regulation by thyroid hormone is absent.<sup>(131)</sup> TR  $\beta 2$  is the key isoform responsible for T3-mediated feedback regulation by hypophysiotropic TRH neurons<sup>(132)</sup>. Targeted disruption of TR  $\beta 2$  expression results in increased TRH mRNA expression in PVN, similar to that found in hypothyroidism. At difference with the anterior pituitary, where ablation of TR  $\beta 2$  or the entire TR  $\beta$  allele produces only a partial TH resistance,<sup>(38)</sup> the lack of TR  $\beta$  is associated with a complete resistance of the modulation of TRH synthesis exerted by severe hypo or hyperthyroidism.<sup>(132)</sup> The physiological source of the T3 causing down regulation of TRH mRNA in the hypothalamus is the subject of ongoing investigations. Somewhat surprisingly, the PVN does not contain the Type 2 5' iodothyronine deiodinase (D2) which is thought to be the source of at least 80% of the intracellular T3 in the central nervous system<sup>(38)</sup> (see Chapter 3). However, studies with T3 containing mini-pumps implanted into thyroidectomized rats indicate that, for normalization of circulating TSH and hypothalamic pre-pro-TRH mRNA, T3 concentrations about twice normal have to be maintained in rat plasma.<sup>(128)</sup> Thus, for both systems (TRH and TSH), feedback regulation requires a source of T3 in addition to that provided by the ambient levels of this hormone. While this T3 seems likely to be produced locally from T4, the main anatomical location of such a process has been identified only recently in the specialized ependymal cells called tanycytes lining the floor and the infralateral wall of the third ventricle between the rostral and caudal poles of the median eminence and the infundibulare recess.<sup>(38)</sup> Tanycytes are one of the major sources of D2, with D2 mRNA expressed in the cell bodies, in the processes and in the end feet<sup>(38)</sup>. Originally believed to only serve as part of the blood-brain barrier, tanycytes have complex functions including active role in endocrine regulation. In particular, T3 locally produced by tanycytes from circulating T4 represents the primary source of T3 involved in the feed-back regulation of hypophysiotropic neurons, unable to express D2.<sup>(38)</sup> The anatomical location of tanycytes places them in a strategic position to extract T4 from the bloodstream or from cerebrospinal fluid after T4 has traversed the choroid plexus (see later, Fig. 4.12). Despite their lipophylic nature, the transport of thyroid hormone into the cells require an active processes involving a long list of transporters.<sup>(135)</sup> Two transporter families have been shown to be important in the transport of thyroid hormones in the brain: the organic anion transporting polypeptide (OATP) and the monocarboxylate transporter (MCT). While the function of OATP is still poorly understood, several lines of evidence support an important role of MCT8, a member of MTC family in central nervous system thyroid hormone transport expressed primarily in neurons and in tanycytes. Several data from both MCT8 KO mice and from humans with MCT8 mutations indicate that lack of functional MCT8 result in hypothyroid TRH neurons, in spite of high circulating T3 concentration, suggesting that MCT8 is necessary of the physiological feed-back regulation.<sup>(38)</sup>

-192	<b>GGCCAGTGCC</b>	TCCGCGCCCC	GGCTCCGG <b>T</b> C	<u>CCCACCGTCC</u>	CCGCCCCAGA
-142	TTTCCGGAGG	AGCAGGCGGG	CGGGGTCCCG	CGGGGCCGGC	<u>TGCCGTCA</u> GC
-92	GCCCCTTCCC	GGCGGCCGCG	ACCCCTCCCC	<u>GC</u> <b>TGACCTCA</b>	CTCGAGCCGC
-42	CGCCTGGCGC	AGATATAAGC	GGCGGCCCAT	CTGAAGAGGG	CTCGGCAGGC
+9	GCCCAGGGTC	CTCAGCGCTG	CAGACTCC <b>TG</b>	<u>ACCT</u> GCCGAC	TGCGGATCCC

Figure 4- 12 The 5' flanking sequence of the human preproTRH gene between -192 and +58 bp. Four potential thyroid response element (TRE, boxed) and two potential CREB binding elements (CRE, underlined) are shown. One sequence (from -60 to -53 bp) has overlapping TRE/CRE bases (bold). (Modified from Wilber & Xu (131))

The synthesis of TRH is under complex transcriptional control sharing several mechanisms besides the negative regulation by thyroid hormone with the TSH  $\beta$  gene. The human TRH gene (Fig 4.10) is located on chromosome 3 (3q13.3 à q21)<sup>(136)</sup>; the 5' flanking sequence of the TRH gene has potential glucocorticoid and cyclic AMP response elements (GRE and CRE).<sup>(114)</sup> There are also potential negative TREs located in this portion of the gene which offer regulatory sites for thyroid hormone control of TRH gene transcription. The thyroid hormone negative regulatory elements of TRH gene are localized in its 5' flanking element (-242 to +54 bp). Four sequences within this region exhibit high degree of homology with consensus sequences for TRE half-sites (AGGTCA) and two of them show also homology with elements implicated in negative regulation by thyroid hormone of TSH  $\beta$  gene.

<sup>(131)</sup> In the absence of thyroid hormone, proTRH gene expression as well as prohormone convertase enzymes (PC1/3 and PC 2) are increased in PVN, while the content of TRH in the median eminence is decreased due to increased secretion of the mature hormone in the portal circulation.<sup>(38)</sup> In contrast, hyperthyroidism is associated with decreased proTRH-mRNA in PVN.<sup>(38)</sup> The negative feed-back of thyroid hormones is exerted directly on hypophysiotropic TRH neurons of PVN which express all isoforms of TRH. The recent availability of transgenic mice lacking either TRH, TR-  $\beta$  isoforms, or both provided evidence for a pivotal role of TRH in the physiological TH feed-back on the hypothalamic pituitary-thyroid (HPT) axis.<sup>(137)</sup> Double TSH and TR-  $\beta$  knockout mice had reduced TH and TSH levels associated with low TSH content in pituitary thyrotrophs and both serum TSH and pituitary TSH content was increased by chronic exogenous TRH administration.<sup>(38)</sup> Thus, the TRH neuron appears to be required for both TSH and TH synthesis and is the predominant locus of the HPT axis.<sup>(137)</sup> However, studies carried out with different animal models of congenital hypothyroidism show that the thyrotrophs exhibit hyperplasia and hypertrophy along with increased TSH mRNA expression not only in the athyreotic Pax8<sup>-/-</sup> mice, but also in TRHR1<sup>-/-</sup> Pax8<sup>-/-</sup> double-knockout mice, which miss a functional thyroid gland and the TRH receptor at pituitary level, suggesting that the stimulation of thyrotrophs proliferation and TSH synthesis is rather a direct consequence of the athyroidism of the animals.<sup>(139)</sup> Further studies are therefore needed to will be required to determine the relative contributions of TRH and TH for bioactive pituitary TSH release.




Figure 4-10 (new) Genomic and promoter structure of TRH. The murine, rat and human TRH genes are composed of three exons and two introns (A). The coding sequence for the precursor protein is present on exons 2 and 3. As depicted, the TRH promoter region precedes the transcription start site in exon 1. The proximal 250-bp sequences of the human, mouse and rat promoters are similar and share the indicated transcription factor binding sites. The location of the CREB binding site (Site 4) and sequences in human (H), mouse (M) and rat (R) are shown. (B,C) Hypothesized schematic representation of the interaction between PCREB and the thyroid hormone receptor at Site 4. (B) Illustrates that in the presence of abundant PCREB, there may be less availability for binding of the thyroid hormone receptor/T3 complex, hence, an increase in TRH gene transcription. When PCREB concentrations fall as shown in (C), increased binding of the thyroid hormone receptor/T3 complex reduces TRH gene transcription (From (112) with permission)

As shown in Fig 4.10, the TRH gene promoter contains potential binding sites for cAMP response element (CRE) binding protein (CREB), and both human and rat TRH genes are positively regulated by cAMP.<sup>(131)</sup> One of the potential CREs of TRH promoter is a sequence that has overlapping TRE/CRE bases -53 to -60 bp (TGACCTCA) (Fig 4-11)<sup>(131)</sup>. There is evidence for competitive interactions of TR $\beta$ 1 molecule and CREB at the overlapping TRE/CRE in the TRH promoter.<sup>(131)</sup> Constructs of TRH promoter with mutations in this overlapping site prevent both the inhibition by TR-T3 complex and the paradoxical stimulation by unliganded TR, underlining the relative importance of TRE/CRE site in relation to the other TRES in the TRH promoter.<sup>(131)</sup>




Figure 4.11 (new) Schematic illustration of the feedback system regulating the hypothalamic-pituitary-thyroid axis. Thyroid hormones exert negative feedback effect at the level of the pituitary and hypophysiotropic TRH neurons. The central feedback effect of thyroid hormones primarily depends on the circulating T4 levels. In the hypothalamus, T4 is converted to T3 by D2 in tanycytes. By volume transmission, T3 secreted from tanycytes reaches the hypophysiotropic TRH neurons, where T3 inhibits the proTRH gene expression via TR- $\beta$ 2 receptors. The set point of the feedback regulation can be altered by two mechanisms: (1) regulation of D2 activity in tanycytes may alter the hypothalamic T3 availability independently from the peripheral T4 concentration. (2) Neuronal afferents can alter the PCREB concentration in the hypophysiotropic TRH neurons that can change the set point of feedback regulation through competition of PCREB and thyroid hormone receptors for the multifunctional binding site (Site 4) of the TRH promoter. ARC, hypothalamic arcuate nucleus; C1-3, C1-3 adrenergic area of the brainstem; CSF, cerebrospinal fluid; DMN, hypothalamic dorsomedial nucleus; ME, median eminence; NTS, nucleus tractus solitarius; PVN, hypothalamic paraventricular nucleus; py, pyramidal tract; sp5, spinal trigeminal tract. (From Fekete & Lechan, 2007, (112) with permission)

A glucocorticoid-responsive element (GRE) is also present in TRH gene promoter<sup>(114)</sup> and glucocorticoid receptor has been identified on TRH neurones of PVN.<sup>(140)</sup> The role of corticosteroids in TRH gene expression is unclear, since both inhibitory and stimulatory effects have been reported.<sup>(38)</sup> The direct effect of glucocorticoids on TRH gene expression is generally stimulatory *in vitro*, but

*in vivo* this activity may be overridden by the complex neuroendocrine reactions following glucocorticoid excess or deficiency. <sup>(141)</sup>

### **TRH interaction with pituitary thyrotrophs and with thyroid hormone**

Although TRH (either maternal or embryonic) is not required for the normal development of the fetal pituitary thyrotrophs and TRH-deficient mice are not hypothyroid at birth, TRH is required later for the postnatal maintenance of the normal thyrotrophs function. <sup>(143)</sup> TRH exerts its activity binding to a specific receptor in the plasma membrane of the thyrotroph to induce the release of TSH and to stimulate TSH synthesis. The TRH receptor of several animal species (including humans) has been cloned and has been identified as a G-protein-coupled receptor with seven highly conserved transmembrane domains. <sup>(38)</sup> Inactivating mutations in the 5'-part of TRH receptor gene are responsible of congenital central hypothyroidism. <sup>(38)</sup> TRH-receptor number and mRNA is increased by glucocorticoids and decreased by thyroid hormone as well as by TRH itself. <sup>(38)</sup> The second messenger for induction of the thyrotroph response to TRH is intracellular Ca<sup>2+</sup> ([Ca<sup>2+</sup>]i). <sup>(38)</sup> TRH was previously believed to act also through stimulation of adenyl cyclase-cAMP pathway, <sup>(104)</sup> but this mechanism was not confirmed by studies carried out with recombinant TRH-receptor transfected in different cell systems. <sup>(154)</sup> TRH activates a complex [Ca<sup>2+</sup>]i response pattern dependent upon both agonist concentration and cell context. The first phase of the TRH response is an acute increase of [Ca<sup>2+</sup>]i within the thyrotrophs via release from internal stores. This is the consequence of increased inositol triphosphate from hydrolysis of phosphatidyl inositol (PI) in the cell membrane. <sup>(38)</sup> The hydrolysis of PI is mediated by G protein activation of phospholipase C and also generates diacylglycerol, which in turn activates intracellular protein kinase C (PKC). Stimulation of extracellular calcium influx through verapamil sensitive channels is also observed after TRH stimulation. <sup>(38)</sup> Both TRH and increased [Ca<sup>2+</sup>]i stimulate intracellular calcium efflux, which helps in terminating the agonist activity. <sup>(38)</sup> In transfection systems in which the TSH β gene promoter has been linked to a reporter gene, both calcium ionophore ionomycin and phorbol esters (a protein kinase C activator) stimulate TSH gene transcription, confirming the key role of these second messengers in mediating TRH activity. <sup>(159)</sup> Both increased [Ca<sup>2+</sup>]i and PKC appear to be independently operative in normal thyrotrophs. <sup>(160)</sup>

The molecular mechanism(s) underlying the stimulation of TSH β gene expression by TRH have been partially elucidated. In GH3 transfected with hTSH β promoter, two distinct regions of human TSH β gene positively responding to TRH were identified between -130 and +37 bp of the gene. <sup>(38)</sup> (Fig. 4-12) The 3'-region corresponds to eight bp of the first exon; the 5'-region ranged between -128 to -60 bp of the 5'-flanking region. <sup>(38)</sup>

### **Inactivation of TRH**

TRH is rapidly inactivated within the central nervous system by a cell-surface peptidase called TRH-degrading ectoenzyme (TRH-DE) <sup>(164)</sup>. TRH-DE is very specific, since there is no other ectopeptidase known capable of degrading TRH and TRH is the only known substrate of this unique enzyme <sup>(164)</sup>. TRH-DE has been purified to homogeneity and cDNA encoding rat TRH-DE has been cloned. In rodents, pituitary TRH-DE mRNA and enzymatic activity are stringently positively regulated by thyroid hormones, and reduced by estrogens. <sup>(164)</sup> This suggests that TRH-DE may act as a regulatory element modulating pituitary TSH secretion. The expression of TRH-DE in brain is high and displays a

distinct distribution pattern, but it is not influenced by peripheral hormones, supporting the concept that brain TRH-DE may act as a terminator of TRH signals.<sup>(164)</sup>

### **Summary of the main steps involved in the hypothalamic-pituitary-thyroid (HPT) axis**

An attempt to summarize the main steps involved in the feedback regulation of the HPT axis is reported in Fig 4.12.<sup>(38)</sup> Thyroid hormones inhibit the effects of TRH on TSH release without interfering with TRH binding to its receptors, but exerting complex negative transcriptional and post-transcriptional activities on TSH synthesis and secretion discussed before. Several factors other thyroid hormones involved in the fine regulation of HPT axis are also depicted in Fig. 4.13 and described in more detail in the following paragraph.




Figure 4- 13 The regulatory region of human TSH- $\beta$  gene (see text for details)

### **Other factors involved in regulation of TSH/TRH synthesis and secretion**

A number of other substances, including ubiquitary and pituitary or thyrotroph-specific transcription factors, hormones, neuropeptides and cytokines influence TSH synthesis and secretion (Table 4-3, Fig. 4-12 and 4-13).

Table 4-3 Predominant Effects of Various Agents on TSH Secretion

STIMULATORY	INHIBITORY
Thyrotropin-releasing hormone (TRH)	Thyroid hormones and analogues
Prostaglandins (?)	Dopamine
$\alpha$ -adrenergic agonists (? Via TRH)	Gastrin
Opioids (humans)	Opioids (rat)
Arginin-vasopressin (AVP)	Glucocorticoids (in vivo)

Glucagon-like peptide 1 (GLP-1)	Serotonin
Galanin	Cholecystokinin (CCK)
Leptin	Gastrin-releasing peptide (GRP)
Glucocorticoids (in vitro)	Vasopressin (AVP)
	Neuropeptide Y (NPY )
	Interleukin 1 $\beta$ and 6
	Tumor necrosis factor $\alpha$

### **Role of Pit-1 and its splicing variants in the regulation of TSH $\beta$ gene expression**

Sequence analysis of the hTSH  $\beta$  promoter reveals three areas with high (75-80%) homology to the consensus sequence for the pituitary-specific transcription factor Pit-1.<sup>(38)</sup> These areas are localized between -128 and -58 bp of the 5'-flanking region. Selective mutation analysis revealed that the integrity of these areas was needed for the stimulatory effect of either TRH or forskolin.<sup>(166)</sup>

Expression of an inactive mutant of Pit-1 decreases TRH stimulation of hTSH  $\beta$ <sup>(38)</sup> and transfection of Pit-1 in cell lines lacking this factor restores cAMP induction of hTSH  $\beta$  gene.<sup>(38)</sup> Taken together these results strongly support an important role of Pit-1 in the regulation of hTSH  $\beta$  gene expression.

Phosphorilation markedly increases the stimulatory activity of Pit-1 in TSH  $\beta$  gene expression,<sup>(166)</sup> and TRH stimulates transient phosphorilation of Pit-1 in GH3 pituitary cells.<sup>(167)</sup>

Further support for a role of Pit-1 in the regulation of TSH  $\beta$  gene expression derives from animal models (dwarf mice) and from clinical syndromes of combined pituitary hormone deficiency (CPHD).<sup>(38)</sup>

Snell and Jackson dwarf mice lack of functioning Pit-1 protein due to a point mutation and a gross structural rearrangement in the Pit-1 gene, respectively.<sup>(38)</sup> Both species show low serum concentration of GH, prolactin and TSH associated to the loss of somato-, lacto- and thyrotropic pituitary cells. Several Pit-1 point mutations and a deletion of the entire coding sequence have been described in patients with CPHD: the effects on TSH secretion differ with the localization of the mutation, but generally result in central hypothyroidism.<sup>(38)</sup>

Although important, the role of Pit-1 for cell-specific expression of TSH  $\beta$  is not as clear as with the GH and PRL genes.<sup>(38)</sup> Attention has been focused on thyrotropin-specific transcription factors, including Pit-1 splicing variants. Of those, a variant called Pit-1T (containing a 14 aminoacid insertion in the transactivation domain) is found only in thyrotropic cells expressing TSH  $\beta$  and increases TSH  $\beta$  promoter activity when transfected in non-thyrotropic cells expressing wild type Pit-1.<sup>(38)</sup> These results suggest that the combination of both Pit-1 and Pit-1T may have a synergistic stimulatory effect of TSH  $\beta$  promoter.<sup>(180)</sup>

### **Other transcription factors involved in TSH $\beta$ gene expression**

As stated before, the transcription factor AP-1 may be involved in modulating the thyroid hormone regulation of TSH- $\beta$  gene expression. Accordingly, a potential AP-1 binding site is present between -1 to +6 bp of the TSH- $\beta$  gene,<sup>(161)</sup> and the integrity of this site is required for maximal stimulation of hTSH- $\beta$  gene.<sup>(181)</sup> Haugen et al.<sup>(182)</sup> described a new 50 kd thyrotroph-specific protein whose binding together with Pit-1 was needed for optimal basal expression of mouse TSH- $\beta$  gene protein which was subsequently identified as the transcription factor GATA-2.<sup>(183)</sup> GATA-2 synergistically with Pit-1 stimulates mouse TSH- $\beta$  promoter activity and is needed for optimal TSH  $\beta$  gene basal activity.

Another pituitary-specific protein (P-Lim), which binds and activates common glycoprotein hormone  $\alpha$  subunit promoter, also synergizes with Pit-1 in the transcriptional activation of TSH  $\beta$  genes in mice. (38) Recently, a syndrome of CPHD including central hypothyroidism has been described in family members carrying mutations of the pituitary transcription factor prophet of pit-1 (PROP-1). (38) Similarly to Pit-1, this finding suggests an important role of PROP-1 for the cell-specific expression of TSH- $\beta$  gene.

## cAMP

Increase of intracellular cAMP stimulate expression of both common  $\alpha$  and TSH  $\beta$  subunit genes. (163) At difference with the TRH gene, this action is probably exerted not through a direct binding to a CRE sequence, but promoting Pit-1 phosphorylation with subsequent activation of TSH- $\beta$  promoter. (38)

## Steroid hormones

Steroid hormones including corticosteroids, estrogen and testosterone modulate TSH  $\beta$  gene expression. Dexamethasone in pharmacological doses decreases serum TSH concentrations of TSH in normal subjects (187) and in patients (188) or rats (189) with TSH-secreting pituitary adenomas, but does not significantly change TSH subunits mRNA levels. (189) This suggests that glucocorticoids may act on TSH biosynthesis at a translational or post-translational level. Furthermore, as discussed before for TRH gene, several other neuroendocrine mechanisms may participate *in vivo* in the modulation of TSH synthesis/secretion by glucocorticoids. In keeping with this concept, it has been recently shown in humans that enhanced hypothalamic somatostatinergic and dopaminergic inhibitory activities are involved in the glucocorticoid-dependent blunting of TSH response to TRH. (190)

Estrogens and testosterone have scanty direct effects on TSH synthesis and secretion in humans. Estrogens mildly reduce  $\alpha$  and  $\beta$  TSH subunit mRNA in hypothyroid rats, (191) perhaps interacting with the same response elements involved in thyroid hormone regulation. Testosterone has similar effects, at least in part explained by its peripheral conversion to estrogen. (192)

## Other hormones, neuropeptides and cytokines

Somatostatin, the major physiological inhibitor of GH secretion, is also an inhibitor of TSH secretion in rats and humans. (38) The physiological relevance of this inhibition is suggested by studies carried out with antibodies to somatostatin whose administration in rats increases serum TSH in basal conditions and after TRH or cold-exposure. (196) Indirect evidence for a physiological role of somatostatin in the regulation of TSH secretion has been obtained in humans by the demonstration that stimulation of the endogenous somatostatin tone by oral glucose inhibits TSH response to TRH. (197) The TSH-inhibiting activity of somatostatin is an acute phenomenon, while long-term treatment with somatostatin analogues does not cause hypothyroidism in man, (38) presumably because the effects of the initial decrease in serum thyroid hormone concentration overrides the inhibitory effects of somatostatin. Somatostatin binds to five distinct types of receptors expressed in the anterior pituitary and brain and differing in binding specificities, molecular weight and linkage to adenyl cyclase. (200) Binding of somatostatin to its receptor causes activation of Gi proteins which in turn inhibit adenylate cyclase. Somatostatin also induces cellular hyperpolarization via modulation of voltage-dependent potassium channels: (201) this mechanism is cAMP-independent and leads to a fall of  $[Ca^{2+}]_i$  by reducing extracellular calcium influx. (202)

In animal models, TSH secretion is affected by other hypothalamic hormones: in particular, corticotropin-releasing hormone (CRH) stimulates TSH secretion in chicken<sup>(203)</sup> through interaction with CRH-receptor-2<sup>(204)</sup> and Melanin-concentrating hormone (MCH) suppresses *in vivo* and *in vitro* TSH release in rats.<sup>(205)</sup>

Neurotransmitters are important direct and indirect modulators in TSH synthesis and secretion. A complex network of neurotransmitters neurons terminates on cells bodies of hypophysiotropic neurons and several neurotransmitters (such as dopamine) are directly released into hypophysial portal blood exerting direct effects on anterior pituitary cells. Furthermore, many dopaminergic, serotoninergic, histaminergic, catecolaminergic, opioidergic and GABAergic systems project from other hypothalamic/brain regions to the hypophysiotropic neurons involved in TSH regulation. These projections are important for a normal TSH circadian rhythm, response to stress and cold exposure, while basal TSH secretion is mainly regulated by intrinsic hypothalamic activity.<sup>(206)</sup> In spite of the difficulty to precisely identify the relative contributions of different neurotransmitter systems in the regulation of TSH secretion, the role of some of them (particularly dopamine and catecholamines) has been rather well defined.

Dopamine, acting via DA2 class of dopamine receptors, inhibits TSH synthesis and release; similarly to somatostatin, this activity is exerted through a decrease in adenylate cyclase.<sup>(38)</sup> Dopamine also inhibits  $\alpha$  and TSH- $\beta$  subunit mRNA and gene transcription in cultured rat anterior pituitary cells.<sup>(63)</sup> In contrast with its inhibitory activity at the thyrotrophs level, dopamine at the hypothalamic levels stimulates both TRH and somatostatin release,<sup>(38)</sup> with opposite effect on TSH secretion.

In contrast to dopamine, adrenergic activation exerts a positive regulation on TSH secretion. Central stimulation of  $\alpha$ -adrenergic pathways increases TSH release in rat, presumably through stimulation of TRH secretion; furthermore,  $\alpha$ 1 adrenergic agonists also enhance TSH release from pituitary cells *in vitro* by mechanisms which are independent of those activated by TRH.<sup>(38)</sup> It is thought that  $\alpha$ -adrenergic activity on thyrotrophs is linked to adenylate cyclase activation since agents increasing intracellular cyclic AMP in these cells can increase TSH release.<sup>(38)</sup>

Opioids inhibit TSH secretion in rats and this action is blocked by the antagonist naloxone,<sup>(216)</sup> while in humans appear to exert a stimulatory effect, especially on nocturnal TSH surge.<sup>(217)</sup> Several other neuropeptides may affect TSH secretion *in vivo* or *in vitro*. Colecistokinin (CCK),<sup>(218)</sup> gastrin-releasing peptide (GRP)<sup>(219)</sup> and neuropeptide Y (NPY)<sup>(38)</sup> exert inhibitory effects, while arginin-vasopressin (AVP),<sup>(221)</sup> glucagon-like peptide-1 (GLP-1)<sup>(222)</sup>, galanin<sup>(223)</sup> and leptin<sup>(38)</sup> stimulate TSH secretion. Although the precise physiological role of these peptides remains to be clarified, it has been recently suggested that they may be important in connecting the nutrition status and thyroid function,<sup>(226)</sup> as discussed in more detail later.

Cytokines have recently been demonstrated to have important effects on TRH or TSH release. Both interleukin 1  $\beta$  (IL-1 $\beta$ ) and tumor necrosis factor  $\alpha$  (cachectin) inhibit TSH basal release,<sup>(38)</sup> while no inhibition is observed on TSH response to TRH.<sup>(231)</sup>, and this effect is independent from thyroid hormone uptake or receptor occupancy. At the same time, IL-1 $\beta$  stimulates the release of corticotropin-releasing hormone and activates the hypothalamic-pituitary-adrenal axis.<sup>(232)</sup> Interleukin-1 $\beta$  is produced in rat thyrotrophs, and this production is markedly increased by bacterial lipopolysaccharide<sup>(38)</sup>. It could thus reduce TSH secretion by either autocrine or paracrine mechanisms. The IL-1 $\beta$ -dependent cytokine interleukin 6 (IL-6) exerts similar inhibitory effects on TSH secretion. Both IL-1 $\beta$  and IL-6 acutely inhibit TSH release from the thyrotrophs, while IL-1 $\beta$  (but not IL-6) also decreases

hypothalamic TRH mRNA and gene expression.<sup>(38)</sup> Both IL-1 $\beta$  and IL-6 stimulate 5'-deiodinase activity in cultured pituitary cells,<sup>(38)</sup> suggesting that increased intrapituitary T4 à T3 conversion may be involved in the inhibitory activity on TSH production. IL-6 is produced by the folliculo-stellate cells of the anterior pituitary<sup>(38)</sup> and, like IL-1 $\beta$ , may regulate TSH release in a paracrine fashion.<sup>(38)</sup> As discussed later, increased concentration of circulating pro-inflammatory cytokines are involved in the alterations of hypothalamic-pituitary-thyroid axis observed in non-thyroidal illnesses.

Quite recently, SIRT1, a NAD-dependent deacetylase, has been proven to be important for TSH secretion by thyrotrophic cells by the pathway SIRT1-phosphatidylinositol-4-phosphate 5-kinase-gamma.<sup>(38)</sup>

Thus an intricate set of relationships within and outside the central nervous system controls the TRH-producing neurones in the medial basal hypothalamus. Alterations in any of these mechanisms can influence TRH and consequently TSH release (Fig 4-13 and 4-14). The relative importance in human physiology of these neural pathways, which have been directly studied only in animal models, is unknown.




Figure 4- 14 Schematic representation of the main factors interacting in the regulation of TSH synthesis and secretion (DA: dopamine; SS: somatostatin;  $\alpha$ -AD:  $\alpha$  adrenergic pathways). Red arrows: stimulation; blue blunted arrows: inhibition

### Short and ultra short-loop feedback control of TSH secretion

In addition to the classic negative feed-back of thyroid hormone on TSH and TRH secretion detailed in the above paragraph, evidence is accumulating that pituitary TSH is able to inhibit TRH secretion at hypothalamic level (short feedback) and TSH secretion at pituitary level (ultra short feedback).<sup>(242)</sup> Early observations of inhibition of TSH secretion by injection of pituitary extracts have been recently corroborated by the demonstration of TSH receptor (together with other pituitary hormone receptors) in the hypothalamus<sup>(38)</sup> and in the folliculo-stellate cells of the adenohypophysis.<sup>(245)</sup> The precise physiological role of short and ultra-short feedback in controlling TRH/TSH secretion remains to be

elucidated. It may be speculated that they concur in the fine tuning of the homeostatic control and in the generation of the pulsatility of TSH secretion. The possibility that thyroid-stimulating autoantibody of Graves' disease recognize hypothalamic and pituitary TSH receptors has also been suggested to explain suppressed serum TSH levels in some euthyroid Graves' patients. <sup>(242)</sup>

## PHYSIOLOGICAL REGULATION OF TSH SECRETION IN HUMANS

A number of experimental paradigms have been used to mimic clinical situations that affect the hypothalamic-pituitary thyroid axis in man. However, with the exception of the studies of thyroid status and iodine deficiency, such perturbations have limited application to humans due to differences in the more subtle aspects of TSH regulation between species. For example, starvation is a severe stress and markedly reduces TSH secretion in rats, but only marginally in humans. Cold stress increases TSH release in the adult rat by  $\alpha$ -adrenergic stimulation, while this phenomenon is usually not observed in the adult human. Thus, it is more relevant to evaluate the consequences of various pathophysiological influences on TSH concentrations in humans rather than to extrapolate from results in experimental animals. This approach has the disadvantage that, in many cases, the precise mechanism responsible for the alteration in TSH secretion cannot be identified. This deficit is offset by the enhanced relevance of the human studies for understanding clinical pathophysiology.

### Normal Physiology

The concentration of TSH can now be measured with exquisite sensitivity using immunometric techniques (see below). In euthyroid humans, this concentration is 0.4-0.5 to 4.0-5.0 mU/L. This normal range is to some extent method-dependent in that the various assays use reference preparations of slightly varying biological potency. There is no crystalline human TSH preparation, so it is not possible to provide a precise molar equivalent for TSH concentrations. Recently, a narrower range (0.5-2.5 mU/L) has beeen proposed in order to exclude subjects with minimal thyroid dysfunction, particularly subclinical hypothyroidism <sup>(246)</sup>, but the issue is still controversial. <sup>(247)</sup> Moreover, data form wide epidemiological studies mostly carried out in iodine sufficient countries like U.S.A., suggest that age together with racial and ethnic factor may significantly affect the respective "normal" TSH range, with higher levels for older caucasian subjects.<sup>(38)</sup> These data differ form the findings previously reported in selected small series of healthy elderly subjects <sup>(250)</sup> suggesting an age-associated trend to lower serum TSH concentrations (see below). The reason(s) for such discrepancy is (are) still not understood. Independently from the "true" normal range of serum TSH, there is substantial evidence that this is genetically controlled, the heritability being estimated between 40-65%. <sup>(251)</sup> As reported in Table 4.4, polymorphisms of several genes encoding potentially involved in the control of HPT axis show a significant link with serum TSH concentrations <sup>(252)</sup> and PDE8B, a gene encoding a high-affinity phosphodiesterase catalyzing the hydrolysis and inactivation of cAMP, has been shown by genome-wide association study to be one of the most important <sup>.(38)</sup>

Table 4.4 Common polymorphisms related to serum thyroid hormones and TSH variation <sup>(252)</sup>

Gene	Polymorphism	Effect on serum					
		T4/T3	T3	rT3	T3/rT3	=	=
TS	T4						
TSHR	Asp727Glu	↓	=	=	=	=	=

	rs10149689 A/ <b>G</b> *	↑	=	=	=	=	=
	rs12050077 A G	↑	=	=	=	=	=
<i>DIO1</i>	D1a- <b>C</b> /T	=	↓	↑	↓		
	D1b-A/ <b>G</b>	=	↑	↓	↑		
	rs2235544 C /A	=	↑	↓	↑		
<i>DIO2</i>	D2-ORFa-Asp3	=	↑ <sup>1</sup>	=	=	=	=
	Thr92Ala	=	=	=	=	=	<sup>2</sup>
	rs225014 C /T	=	=	=	=	=	<sup>3</sup>
<i>THRB</i>	TRHB-in9. A /G	(↑)	=	=	=	=	=
<i>PDE8B</i>	rs4704397 A /G	↑	=	=	=	=	=

\* Alleles associated with the specified trait are reported in **bold**<sup>1</sup>Only in young subjects<sup>2</sup> Influence L-T4 dose needed to normalize serum TSH in hypothyroid patients

<sup>3</sup> Influence psychological well-being of hypothyroid patients on L-T4 therapy

Free α subunit is also detectable in serum with a normal range of 1 to 5 µg/L, but free TSH β is not detectable<sup>(38)</sup>. Both the intact TSH molecule and the α subunit increase in response to TRH. The α subunit is also increased in post-menopausal women; thus the level of gonadal steroid production needs to be taken into account in evaluating α subunit concentrations in women. In most patients with hyperthyroidism due to TSH-producing thyrotroph tumors, there is an elevation in the ratio of α subunit to total TSH.<sup>(38)</sup> In the presence of normal gonadotropin this ratio is calculated by assuming a molecular weight for TSH of 28,000 and for α subunit of 13,600 Da. The approximate specific activity of TSH is 0.2 mU/mg. To calculate the molar ratio of a subunit to TSH, the concentration of α subunit (in µg/L) is divided by the TSH concentration (in mU/L) and this result multiplied by 10. The normal ratio is <1.0 and it is usually elevated in patients with TSH-producing pituitary tumor but is normal in patients with thyroid hormone resistance unless they are post menopausal.

The volume of distribution of TSH in humans is slightly larger than the plasma volume, the half-life is about 1 hour, and the daily TSH turnover between 40 and 150 mU/day.<sup>(257)</sup> Patients with primary hypothyroidism have serum TSH concentrations greater than 5 and up to several hundred mU/L.<sup>(103)</sup> In patients with hyperthyroidism due to Graves' disease or autonomous thyroid nodules, TSH is suppressed with levels which are inversely proportional to the severity and duration of the hyperthyroidism, down to as low as <0.004 mU/L.<sup>(38)</sup>

TSH secretion in humans is pulsatile.<sup>(38)</sup> The pulse frequency is slightly less than 2 hours and the amplitude approximately 0.6 mU/L. The TSH pulse is significantly synchronized with PRL pulse: this phenomenon is independent from TRH and suggests the existence of unidentified underlying pulse generator(s) for both hormones.<sup>(264)</sup> The frequency and amplitude of pulsations increases during the evening reaching a peak at sleep onset, thus accounting for the circadian variation in basal serum TSH levels.<sup>(38)</sup> The maximal serum TSH is reached between 21:00 and 02:00 hours and the difference between the afternoon nadir and peak TSH concentrations is 1 to 3 mU/L. Sleep prevents the further rise in TSH as reflected in the presence of increases in TSH to 5-10 mU/ml during sleep deprivation.<sup>(38)</sup> The circadian variation of TSH secretion is probably the consequence of a varying dopaminergic tone modulating the pulsatile TSH stimulation by TRH.<sup>(269)</sup> There is little, if any, significant seasonal

change in basal TSH nor are there any gender-related differences in either the amplitude or frequency of the TSH pulses.<sup>(263)</sup> The diurnal rhythmicity of serum TSH concentration is maintained in mild hyper- and hypothyroidism, but it abolished in severe short-term primary hypothyroidism, suggesting that the complete lack of negative feedback to the hypothalamus or pituitary or both may override the central influences on TSH secretion.<sup>(270)</sup>

Age does not have a major effect on serum TSH with the exception of the extremes.<sup>(250)</sup> There is a marked increase in serum TSH in neonates which peaks within the first few hours of delivery returning towards normal over the next few days (see Chapter 15). It is thought to be a consequence of the marked reduction in environmental temperature at birth. Serum TSH concentrations in apparently euthyroid patients over the age of 70 may be somewhat reduced,<sup>(38)</sup> although usually this is a pathological finding indicating either exogenous or endogenous thyrotoxicosis.<sup>(273)</sup> However, slightly decreased serum TSH has been also documented in elderly patients without clinical or subclinical hyperthyroidism, as assessed by 4 year follow-up,<sup>(272)</sup> and in healthy centenarians.<sup>(274)</sup> An age-dependent reduction of daily TSH secretion rate has been reported in humans.<sup>(275)</sup> The physiological nyctohemeral rhythm of TSH is maintained in the elderly, but the nocturnal peak is blunted.<sup>(276)</sup>

## TSH in Pathophysiological States

### Nutrition

In the rat, starvation causes a marked decrease in serum TSH and thyroid hormones. While there is an impairment of T4 to T3 conversion in the rat liver due to a decrease in both thiol co-factor and later in the Type 1 deiodinase,<sup>(38)</sup> the decrease in serum T3 in the fasted rat is primarily due to the decrease in T4 secretion consequent to TSH deficiency.<sup>(38)</sup> In humans, starvation and moderate to severe illness are also associated with a decrease in basal serum TSH, pulse amplitude and nocturnal peak.<sup>(38)</sup> In the acutely-fasted man, serum TSH falls only slightly and TRH responsiveness is maintained, although blunted.<sup>(38)</sup> This suggests that the thyrotroph remains responsive during short-term fasting and that the decrease in TSH is likely due to changes due to decreased TRH release. There is evidence to support this in animal studies, showing reduced TRH gene expression in fasted rats.<sup>(38)</sup> Administration of anti-somatostatin antibodies prevents the starvation induced serum TSH falls in rats, suggesting a role for hypothalamic somatostatinergic pathways.<sup>(291)</sup> However, fasting-induced changes in dopaminergic tone do not seem to be sufficient to explain the TSH changes.<sup>(38)</sup>

Recent studies provide compelling evidence that starvation-induced fall in leptin levels (Fig. 4-14) plays a major role in the decreased TSH and TSH secretion of fasted animals and, possibly, humans.<sup>(38)</sup> This concept stems from the observation that administration of leptin prevents the starvation-induced fall of hypothalamic TRH.<sup>(294)</sup> The mechanisms involved in this phenomenon include decreased direct stimulation by leptin of TRH production by neurons of the PVN,<sup>(38)</sup> as well as indirect effects on distinct leptin-responsive neuroendocrine circuits communicating with TRH neurons.<sup>(38)</sup> The direct stimulatory effects of leptin on TRH production are mediated by binding to leptin receptors, followed by STAT3 activation and subsequent binding to the TRH promoter.<sup>(38)</sup> One of the latter circuits has been identified in the melanocortin pathway, a major target of leptin action. This pathway involves 2 ligands expressed in distinct populations of arcuate nucleus neurons in the hypothalamus (the  $\alpha$ -MSH and the Agouti receptor protein [AgRP]) and the melanocortin 4 receptor (MC4R) on which these ligands converge, but exert antagonistic effects (stimulation by  $\alpha$ -MSH;

inhibition by AgRP). Leptin activates MC4R by increasing the agonist  $\alpha$ -MSH and by decreasing the antagonist AgRP and this activation is crucial for the anorexic effect of leptin. The specific involvement of melanocortin pathway in TRH secretion is suggested by the presence of  $\alpha$ -MSH in nerve terminals innervating hypothalamic TRH neurons in rat<sup>(299)</sup> and human<sup>(300)</sup> brains and by the ability of  $\alpha$ -MSH to stimulate and of AgRP to inhibit hypothalamus-pituitary thyroid axis both in vitro and in vivo.<sup>(38)</sup> The activities of  $\alpha$ -MSH and AgRP on thyroid axis are fully mediated by MCR4, as shown by experiment carried out on MCR4 knock out mice.<sup>(38)</sup> Fasting may inhibit the hypothalamic-pituitary-thyroid axis also via the orexogenic peptide NPY, which inhibits TRH synthesis<sup>194.1</sup> by activation of Y1 and Y5 receptors in hypophysiotropic neurons of the hypothalamic paraventricular nucleus.<sup>(302)</sup> At least two distinct population of NPY neurons innervate hypophysiotropic TRH neurons,<sup>(303)</sup> suggesting that NPY is indeed an important regulator of hypothalamic-pituitary-thyroid axis.

A further contributing cause of the decreased TSH release in fasting may be an abrupt increase in the free fraction of T4 due to the inhibition of hormone binding by free fatty acids.<sup>(304)</sup> This would cause an increase in pituitary T4 and, hence, in pituitary nuclear T3. Fasting causes a decrease in the amplitude of TSH pulses, not in their frequency.<sup>(305)</sup>

Ingestion of food results in an acute decline of serum TSH concentration: this is the consequence of meal composition, rather than stomach distension.<sup>(306)</sup> Long-term overfeeding is associated to a transient increase of serum T3 concentration and a sustained increased response of TSH to TRH.<sup>(307)</sup>

Taken together, the above data provide compelling evidence that hypothalamic-pituitary-thyroid axis is strictly related to the mechanisms involved in weight control. In keeping with this concept, preliminary epidemiological studies suggest that small differences in thyroid function may be important for the body mass index and the occurrence of obesity in the general population.<sup>(308)</sup>

## Illness

The changes in circulating TSH which occur during fasting are more exaggerated during illness. In moderately ill patients, serum TSH may be slightly reduced but the serum free T4 does not fall and is often mildly increased.<sup>(38)</sup> However, if the illness is severe and/or prolonged, serum TSH will decrease and both serum T4 (and of course T3) decrease during the course of the illness (see Chapter 5). This may be due to decreased pulse amplitude and nocturnal TSH secretion.<sup>(38)</sup> Since such changes are short-lived, they do not usually cause symptomatic hypothyroidism. They are often associated with an impaired TSH release after TRH.<sup>(282)</sup> However, the illness-induced reductions in serum T4 and T3 will often be followed by a rebound increase in serum TSH as the patient improves. This may lead to a transient serum TSH elevation in association with the still subnormal levels of circulating thyroid hormones and thus be mistaken for primary hypothyroidism.<sup>(315)</sup> On occasion, transient TSH elevation occurs while the patient is still ill. The pathophysiology of this apparent thyroid gland resistance to TSH is not clear,<sup>(316)</sup> although this phenomenon could be the consequence of reduced TSH bioactivity, possibly consequent to abnormal sialylation.<sup>(317)</sup> The transient nature of these changes is reflected in normalization of the pituitary-thyroid axis after complete recovery. It is currently not clearly established whether the above abnormalities on hypothalamic-pituitary-thyroid axis during critical illness reflect an adaptation of the organism to illness or instead a potentially harmful condition leading to hypothyroidism at tissue level.<sup>(38)</sup>

## **Neuropsychiatric disorders**

Certain neuropsychiatric disorders may also be associated with alterations in TSH secretion. In patients with anorexia nervosa or depressive illness, serum TSH may be reduced and/or TRH-induced TSH release blunted.<sup>(320)</sup> Such patients often have decreases in the evening enhancement of TSH secretion.<sup>(321)</sup> The etiology of these changes is not known though it has been speculated that they are a consequence of abnormal TRH secretion.<sup>(38)</sup> The latter is supported by observations that TRH concentrations in cerebrospinal fluid of some depressed patients are elevated.<sup>(38)</sup> There may be a parallel in such patients between increases in TRH and those of ACTH secretion.<sup>(326)</sup> In agreement with this are the increased serum T4 and TSH levels sometimes found at the time of admission to psychiatric units.<sup>(38)</sup>

## **Mechanisms involved in the hypothalamic-pituitary-thyroid axis suppression in non-thyroidal illnesses**

The precise mechanism(s) underlying the suppression of the hypothalamic-pituitary-thyroid axis are only partially known. Evidence for a direct involvement of TRH-producing neurones in humans has been recently provided by the demonstration of low levels of TRH mRNA in PVN of patients died for nonthyroidal disease.<sup>(328)</sup> Alteration in neuroendocrine pathways including opioidergic, dopaminergic and somatostatinergic activity have been advocated, but in acutely ill patients the major role appears to be played by glucocorticoids.<sup>(329)</sup> (See below for a more detailed discussion). Activation of pro-inflammatory cytokine pathways is another mechanism potentially involved in the suppression of TSH secretion in nonthyroidal illness. As discussed before, IL-2 $\beta$ , TNF $\alpha$  and IL-6 exert *in vivo* and *in vitro* a marked inhibitory activity on TRH-TSH synthesis/secretion. High levels of pro-inflammatory cytokines (particularly IL-6 and TNF- $\alpha$ ) have been described in sera of patients with nonthyroidal illnesses.<sup>(38)</sup> Serum cytokine concentration is directly correlated with the severity of the underlying disease and to the extent of TSH and thyroid hormone abnormalities observed in these patients. Furthermore, cytokines also affect thyroid hormone secretion, transport and metabolism providing all the characteristics to be considered important mediators of thyroid hormone abnormalities observed in nonthyroidal illness.<sup>(38)</sup>

## **Effects of Hormones and Neuropeptides**

### **Dopamine and dopamine agonists**

Dopamine and dopamine agonists inhibit TSH release by mechanisms discussed earlier. Dopamine infusion can overcome the effects of thyroid hormone deficiency in the severely ill patient, suppressing the normally elevated TSH of the patient with primary hypothyroidism nearly into the normal range.<sup>(38)</sup> Dopamine causes a reduction of the amplitude of TSH pulsatile release, but not in its frequency.<sup>(312)</sup> However, chronic administration of dopamine agonists, for example in the treatment of prolactinoma, does not lead to central hypothyroidism despite the fact that there is marked decrease in the size of the pituitary tumor and inhibition of prolactin secretion.

### **Glucocorticoids**

The acute administration of pharmacological quantities of glucocorticoids will transiently suppress TSH.<sup>(38)</sup> The mechanisms responsible for this effect may act both at hypothalamic and pituitary level,

as discussed before. Recently, direct evidence of suppressed TRH synthesis was provided by an autopic study showing reduced hypothalamic TRH mRNA expression in subjects treated with corticosteroids before death.<sup>(340)</sup> TSH secretion recovers and T4 production rates are generally not impaired. In Cushing's syndrome, TSH may be normal or suppressed and, in general, there is a decrease in serum T3 concentrations relative to those of T4.<sup>(338)</sup> High levels of glucocorticoid inhibit basal TSH secretion slightly and may influence the circadian variation in serum TSH.<sup>222</sup> Perhaps as a reflection of this, a modest serum TSH elevation may be present in patients with Addison's disease.<sup>(38)</sup> TSH normalizes with glucocorticoid therapy alone if primary hypothyroidism is not also present. Similarly to patients treated with long-acting somatostatin analogs, patients receiving long-term glucocorticoid therapy do not have sustained reduction of serum TSH nor does hypothyroidism develop, because of the predominant effect of reduced thyroid hormone secretion in stimulating TSH secretion.<sup>(343)</sup>

### Gonadal steroids

Aside from the well described effects of estrogen on the concentration of thyroxine binding globulin (TBG), estrogen and testosterone have only minor influences on thyroid economy (see Chapters 5 and 14). In contrast with the mild inhibitory activity on  $\alpha$  and TSH  $\beta$  gene subunit expression described in rats,<sup>(191)</sup> in humans TSH release after TRH is enhanced by estradiol treatment perhaps because estrogens increase TRH receptor number.<sup>(38)</sup> Treatment with the testosterone analog, fluoxymesterone, causes a significant decrease in the TSH response to TRH in hypogonadal men,<sup>(346)</sup> possibly due to an increase in T4 to T3 conversion by androgen.<sup>(347)</sup> This and the small estrogen effect may account for the lower TSH response to TRH in men than in women although there is no difference in basal TSH levels between the sexes. This is one of the few instances where there is not a close correlation between basal TSH levels and the response to TRH (see below).

### Growth hormone (GH)

The possibility that hypothyroidism could be induced by GH replacement in GH-deficient children was raised in early studies.<sup>(38)</sup> However, these patients received human pituitary GH which in some cases was contaminated with TSH, perhaps inducing TSH antibodies. More recent studies employing recombinant GH have shown no significant changes in TSH concentrations during therapy of adults with GH deficiency.<sup>(350)</sup> Growth hormone does cause an increase in serum free T3, a decrease in free T4, and an increase in the T3 to T4 ratio in both T4-treated and T4 untreated patients. This suggests that the GH-induced increase in IGF-1 stimulates T4 to T3 conversion. In keeping with this concept, IGF-1 administration in healthy subjects is followed by a fall in serum TSH concentration.<sup>(351)</sup>

### Catecholamines

At difference with rat, there is scanty evidence of an adrenergic control of TSH secretion in humans. Acute infusions of  $\alpha$  or  $\beta$  adrenergic blocking agents or agonists for short periods of time do not affect basal TSH,<sup>(38)</sup> although a small stimulatory activity for endogenous adrenergic pathways is suggested by other studies.<sup>(38)</sup> Furthermore, there is no effect of chronic propranolol administration on TSH secretion even though there may be modest inhibition of peripheral T4 to T3 conversion if amounts in excess of 160 mg/day are given.<sup>(356)</sup> Evidence of a tonic inhibition of TSH secretion mediated by endogenous catecholamines has been obtained in women during the early follicular phase of the menstrual cycle.<sup>(357)</sup>

# The Response of TSH to TRH in Humans and the Role of Immunometric TSH Assays

In the last decade, the application of TSH measurements to the evaluation of patients with thyroid disease has undergone a revolutionary change. This is due to the widespread application of the immunometric TSH assay. This assay uses monoclonal antibodies which bind one epitope of TSH and do not interfere with the binding of a second monoclonal or polyclonal antibody to a second epitope. The principle of the test is that TSH serves as the link between an immobilized antibody binding TSH at one epitope and a labelled (radioactive, chemiluminescent or other tag) monoclonal directed against a second portion of the molecule. This approach has improved both sensitivity and specificity by several orders of magnitude. Technical modifications have led to successive “generations” of TSH assays with progressively greater sensitivities.<sup>218,316</sup> The first generation TSH assay is considered to be the standard radioimmunoassay which generally has minimal detection limits of 1-2 mU/L. The “second” generation (first generation immunometric) assay improved the sensitivity to 0.1-0.2 mU/L and the “third” reduced the sensitivity to approximately 0.005 mU/L. Third generation assays are currently being introduced into many clinical laboratories. From a technical point-of-view, the American Thyroid Association recommendations are that third generation assays should be able to quantitative TSH in the 0.010 to 0.020 mU/L range on an interassay basis with a coefficient of variation of 20% or less.<sup>(358)</sup> The most recent development is an assay with a minimal usable sensitivity of 0.0004 mU/L. Such assays are currently available only in specialized laboratories. It would appear that the third generation assays will provide sufficient sensitivity for even the most rigorous clinical applications. As assay sensitivity has improved, the normal range has not changed, remaining between approximately 0.5 and 5.0 mU/L in most laboratories. However, the TSH concentrations in the sera of patients with severe thyrotoxicosis secondary to Graves’ disease have been lower with each successive improvement in the TSH assays: using a fourth generation assay, the serum TSH is <0.004 mU/L in patients with severe hyperthyroidism.<sup>(260)</sup> While the potential for such high sensitivity is inherent to the technology, the clinician should always ascertain that the performance in his/her clinical laboratory meets the appropriate sensitivity criteria before assuming that an assay stipulated to be “second” or “third” generation is achieving that sensitivity on site.<sup>(38)</sup>

The primary consequence of the availability of the sensitive TSH assays is to allow the substitution of a basal TSH measurement for the TRH test in patients suspected of thyrotoxicosis.<sup>(38)</sup> Nonetheless, it is appropriate to review the results of TRH tests from the point-of-view of understanding thyroid pathophysiology, particularly in patients with hyperthyroidism or autonomous thyroid function. In healthy individuals bolus i.v. injection of TRH is promptly followed by a rise of serum TSH concentration peaking after 20 to 30 minutes. The magnitude of the TSH peak is proportional to the logarithm of TRH doses between 6.25 up to  $^3400$   $\mu$ g, is significantly higher in women than in men and declines with age.<sup>(38)</sup> The individual TSH response to TRH is very variable and declines after repeated TRH administrations at short time intervals.<sup>(363)</sup> In the presence of normal TSH bioactivity and adequate thyroid functional reserve, serum T3 and T4 also increase 120 – 180 minutes after TRH injection.<sup>(363)</sup> There is a tight correlation between the basal TSH and the magnitude of the TRH-induced peak TSH. (Fig. 4-12) Using a normal basal TSH range of 0.5 to 5 mU/L, the TRH response 15 to 20 minutes after 500 mg TRH (intravenously) ranges between 2 and 30 mU/L. The lower responses are found in patients with lower (but still normal) basal TSH levels.<sup>(260)</sup> These results are quite consistent with older studies using radioimmunoassays.<sup>(364)</sup> When the TSH response to TRH of all patients (hypo-, hyper- and euthyroid) is analyzed in terms of a “fold” response, the highest response (approximately 20 fold) occurs at a basal TSH of 0.5 mU/L and falls to less than 5 at either markedly

subnormal or markedly elevated basal serum TSH concentrations (Fig. 4-16).<sup>(260)</sup> Thus a low response can have two explanations. The low response in patients with hyperthyroidism and a reduced basal TSH is due to refractoriness to TRH or depletion of pituitary TSH as a consequence of chronic thyroid hormone excess. In patients with primary hypothyroidism, the low fold-response reflects only the lack of sufficient pituitary TSH to achieve the necessary increment over the elevated basal TSH.




Figure 4-16 Relationship between basal and absolute (TRH stimulated-basal TSH) TRH-stimulated TSH response in 1061 ambulatory patients with an intact hypothalamic-pituitary (H-P) axis compared with that in untreated and T4-treated patients with central hypothyroidism. (From C.A. Spencer et al, (260) with permission)

Although, as stated before, the clinical relevance of the TRH test is presently limited, there are still some conditions in which the test may still be useful. These include subclinical primary hypothyroidism, central hypothyroidism, the syndromes of inappropriate TSH secretion and non-thyroidal illnesses.

In patients with normal serum thyroid hormone concentrations and borderline TSH, an exaggerated TSH response to TRH not followed by an adequate increase in serum thyroid hormone levels may confirm the presence of subtle primary hypothyroidism.<sup>(363)</sup>

An abnormal relationship between the basal TSH and the TRH-response is found in patients with central hypothyroidism. Here the fold TSH response to TRH is lower than normal.<sup>(38)</sup> Again, however, TRH testing does not add substantially to the evaluation of such patients in that the diagnosis of central hypothyroidism is established by finding a normal or slightly elevated basal TSH in the presence of a significantly reduced free T4 concentration. While statistically<sup>(260)</sup> lower and sometimes delayed increments in TSH release after TRH infusion are found in patients with pituitary as opposed to hypothalamic hypothyroidism, the overlap in the TSH increments found in patients with these two conditions is sufficiently large<sup>(38)</sup> that other diagnostic technologies, such as MRI, must be used to

provide definitive localization of the lesion in patients with central hypothyroidism . It should be recalled that the TRH test may be useful in the diagnosis and follow-up of several pituitary disorders, but the discussion of this point is beyond the purpose of this chapter.

TRH test still provides fundamental information in the differential diagnosis of hyperthyroidism due to TSH-secreting adenomas from syndromes with non-neoplastic TSH hypersecretion due to pituitary selective or generalized thyroid hormone resistance. In all the above conditions increased or “inappropriately normal” serum TSH concentrations are observed the presence of elevated circulating thyroid hormone levels. However in most (>92%) of TSH-secreting adenomas serum TSH does not increase after TRH, while TRH responsiveness is observed in >95% of patients with nontumoral inappropriate TSH secretion. <sup>(38)</sup>

Perhaps of most interest pathophysiologically is the response to TRH in patients with non-thyroidal illness and either normal or low free T4 indices (Fig. 4-12). Results from these patients fit within the normal distribution in terms of the relationship between basal TSH (whether suppressed or elevated) and the fold-response to TRH. Thus the information provided by a TRH infusion test adds little to that obtained from an accurate basal TSH measurement. <sup>(367)</sup> With respect to the evaluation of sick patients, while basal TSH values are on average higher than in patients with thyrotoxicosis, there is still some overlap between these groups. <sup>(38)</sup> This indicates that even with second or third generation TSH assays, it may not be possible to establish that thyrotoxicosis is present based on a serum TSH measurement in a population which includes severely ill patients.

## **CLINICAL APPLICATION OF TSH MEASUREMENTS AND SUMMARY**

Table 4-5 lists conditions in which basal TSH values may be altered as practical examples of the pathophysiology of the hypothalamic-pituitary thyroid axis. This subject is also discussed in Chapter 6 from the standpoint of clinical diagnosis. This section also serves as a summary of the clinically relevant points in this chapter.

### **Clinical situations associated with subnormal TSH values**

The most common cause of a reduced TSH in a non-hospitalized patient is thyroid hormone excess. This may be due to endogenous hyperthyroidism or excess exogenous thyroid hormone. The degree of suppression of basal TSH is in proportion to the degree and duration of the excess thyroid hormone. The reduced TSH is the pathophysiological manifestation of the activation of the negative feedback loop.

While a low TSH in the presence of elevated thyroid hormones is logical, it results from multiple causes. Prolonged excessive thyroid hormone causes physiological “atrophy” of the thyroid stimulatory limb of the hypothalamic-pituitary thyroid axis. Thus, TRH synthesis is reduced, TRH mRNA in the PVN is absent, TRH receptors in the thyrotroph may be reduced; and the concentration of TSH  $\beta$  and  $\alpha$  subunits and both mRNAs in the thyrotroph are virtually undetectable. Therefore, it is not surprising that several months are usually required for the re-establishment of TSH secretion after the relief of thyrotoxicosis. This is especially well seen in patients with Graves' disease after surgery or radioactive iodine, in whom TSH remains suppressed despite a rapid return to a euthyroid or even hypothyroid functional status. <sup>(38)</sup> Since TRH infusion will not increase TSH release in this situation, it is clear that the thyrotroph is transiently dysfunctional. <sup>(372)</sup> A similar phenomenon occurs after excess thyroid

hormone treatment is terminated, and after the transient hyperthyroidism associated with subacute or some variants of autoimmune thyroiditis, though the period of suppression is shorter under the latter circumstances.<sup>(373)</sup> This cause of reduced circulating thyroid hormones and reduced or normal TSH should be distinguishable from central hypothyroidism by the history.

Severe illness is a common cause of TSH suppression although it is not often confused with thyrotoxicosis. Quantitation of thyroid hormones will generally resolve the issue.<sup>(304)</sup> Patients receiving high-dose glucocorticoids acutely may also have suppressed TSH values although chronic glucocorticoid therapy does not cause sufficient TSH suppression to produce hypothalamic-pituitary hypothyroidism (see above).

Exogenous dopamine suppresses TSH release. Infusion of 5-7.5 mg/Kg/min to normal volunteers causes an approximately 50% reduction in the concentrations of TSH and consequent small decreases in serum T4 and T3 concentrations.<sup>(337)</sup> In critically ill patients, this effect of dopamine can be superimposed on the suppressive effects of acute illness on thyroid function, reducing T4 production to even lower levels.<sup>331</sup> Dopamine is sufficiently potent to suppress TSH to normal levels in sick patients with primary hypothyroidism.<sup>(337)</sup> This needs to be kept in mind when evaluating severely ill patients for this condition. Dopamine antagonists such as metoclopramide or domperidone cause a small increase in TSH in humans. However, somewhat surprisingly, patients receiving the dopamine agonist bromergocryptine, do not become hypothyroid. Although L-dopa causes a statistically significant reduction in the TSH response to TRH, patients receiving this drug also remain euthyroid.<sup>(343)</sup>

Studies in animals have suggested that pharmacological amounts of retinoids may decrease serum TSH concentration (see also paragraph “Effect of Thyroid Hormone on TSH Secretion”).<sup>(38)</sup> Recently, severe central hypothyroidism associated with very low serum TSH concentration has been reported in patients with cutaneous T-cell lymphoma treated with high-dose bexarotene, a retinoid X receptor-selective ligand able to suppress TSH secretion.<sup>(375)</sup>

The capacity of hCG to function as a thyroid stimulator is discussed in Chapter 14. This may be manifested in patients with normal pregnancy as a slightly subnormal TSH during the first trimester (0.2 – 0.4 mU/L) or by frank, though mild, hyperthyroidism in patients with choriocarcinoma or molar pregnancy.<sup>(376)</sup>

Patients with acute psychosis or depression and those with agitated psychoses may have high thyroid hormone levels and suppressed or elevated TSH values. The etiology of the alterations in TSH are not known. Those receiving lithium for bipolar illness may also have elevated TSH values due to impairment of thyroid hormone release. Patients with underlying autoimmune thyroid disease or multinodular goiter are especially susceptible.<sup>(377)</sup> A small fraction of elderly patients, particularly males, have subnormal TSH levels with normal serum thyroid hormone concentrations. It is likely that this reflects mild thyrotoxicosis if it is found to be reduced on repeated determinations.

Congenital central hypothyroidism with low serum TSH may result from mutations affecting TSH  $\beta$  gene or the Pit-1 gene (see paragraphs “The Thyroid-Stimulating Hormone Molecule” and “Role of Pit-1 and its splicing variants in the regulation of TSH  $\beta$  gene expression”).

## Causes of an elevated TSH

Primary hypothyroidism is the most common cause of an elevated serum TSH. The serum free T4 is low normal or reduced in such patients but the serum free T3 values remain normal until the level of thyroid function has markedly deteriorated.<sup>(103)</sup> Another common cause of an elevated TSH in an

iodine-sufficient environment is the transient elevation which occurs during the recovery phase after a severe illness.<sup>(38)</sup> In such patients a “reawakening” of the hypothalamic-pituitary thyroid axis occurs pari passu with the improvement in their clinical state. In general, such patients do not have underlying thyroid dysfunction. Iodine deficiency is not a cause of elevated TSH in Central and North America but may be in certain areas of Western Europe, South America, Africa and Asia.

The remainder of the conditions associated with an elevated TSH are extremely rare. Inherited (autosomal recessive) forms of partial (euthyroid hyperthyrotropinemia) or complete (congenital hypothyroidism) TSH resistance have been recently described associated to inactivating point mutations of the TSH receptor gene.<sup>(38)</sup> Interestingly, inherited dominant forms of partial TSH resistance have also been described in the absence of TSH receptor gene mutations.<sup>(38)</sup> No TSH nor Gsa gene mutations were detected in these exceptional cases.<sup>(380)</sup> and the underlying molecular defect(s) remain(s) to be elucidated. More frequently, in a patient who has an elevated serum FT<sub>4</sub>, the presence of TSH at normal or increased levels should lead to a search for either resistance to thyroid hormone or a thyrotroph tumor. Hypothalamic-pituitary dysfunction may be associated with normal or even modest increases in TSH are explained by the lack of normal TSH glycosylation in the TRH-deficient patient. The diagnosis is generally made by finding a serum free T<sub>4</sub> index which is reduced to a greater extent than expected from the coincident serum TSH. Psychiatric illness may be associated with either elevated or suppressed TSH, but the abnormal values are not usually in the range normally associated with symptomatic thyroid dysfunction. The effect of glucocorticoids to suppress TSH secretion has already been mentioned. This is of relevance in patients with Addison’s disease in whom TSH may be slightly elevated in the absence of primary thyroid disease.

Lastly, while most of the artifacts have been eliminated from the immunometric TSH assays, there remains the theoretical possibility of an elevated value due to the presence of endogenous antimouse gamma globulin antibodies.<sup>(38)</sup> These antibodies, like TSH, can complex the two TSH antibodies resulting in artificially elevated serum TSH assay results in euthyroid patients. Such artifacts can usually be identified by finding non-linear results upon assay of serial dilutions of the suspect serum with that from patients with a suppressed TSH.




Figure 4- 15 Role fasting, somatostatin (SS) pathways and leptin on TRH and TSH secretion. POMC: pro-opiomelanocortin;  $\alpha$ -MSH:  $\alpha$ -melanocyte-stimulating hormone; AGRP: Agouti receptor protein; NPY: neuropeptide Y; MC4R: melanocortin 4 receptor; PVN: paraventricular nucleus Red arrows: stimulation; blue arrows: inhibition; for further details, see text.

## References

1. Shupnik, M.A., Ridgway, E.C.& Chin, W.W. (1989). Molecular biology of thyrotropin. *Endocr Rev* , 10 , 459-475.
2. Grossmann, M., Weintraub, B.D.& Szkudlinski, M.W. (1997). Novel insights into the molecular mechanisms of human thyrotropin action: structural, physiological, and therapeutic implications for the glycoprotein hormone family. *Endocr Rev* , 18 , 476-501.
3. Ridgway, E.C., Weintraub, B.D.& Maloof, F. (1974). Metabolic clearance and production rates of human thyrotropin. *J Clin Invest* , 53 , 895-903.
4. Kourides, I.A., Re, R.N., Weintraub, B.D., Ridgway, E.C.& Maloof, F. (1977). Metabolic clearance and secretion rates of subunits of human thyrotropin. *J Clin Invest* , 59 , 508-516.
5. Naylor, S.L., Chin, W.W., Goodman, H.M., Lalley, P.A., Grzeschik, K.H.& Sakaguchi, A.Y. (1983). Chromosome assignment of genes encoding the alpha and beta subunits of glycoprotein hormones in man and mouse. *Somatic Cell Genet* , 9 , 757-770.
6. Gordon, D.F., Wood, W.M.& Ridgway, E.C. (1988). Organization and nucleotide sequence of the gene encoding the beta-subunit of murine thyrotropin. *DNA* , 7 , 17-26.

7. Burnside, J., Buckland, P.R.& Chin, W.W. (1988). Isolation and characterization of the gene encoding the alpha-subunit of the rat pituitary glycoprotein hormones. *Gene* , 70 , 67-74.
8. Carr, F.E., Need, L.R.& Chin, W.W. (1987). Isolation and characterization of the rat thyrotropin beta-subunit gene. Differential regulation of two transcriptional start sites by thyroid hormone. *J Biol Chem* , 262 , 981-987.
9. Wondisford, F.E., Radovick, S., Moates, J.M., Usala, S.J.& Weintraub, B.D. (1988). Isolation and characterization of the human thyrotropin beta-subunit gene. Differences in gene structure and promoter function from murine species. *J Biol Chem* , 263 , 12538-12542.
10. Tatsumi, K., Hayashizaki, Y., Hiraoka, Y., Miyai, K.& Matsubara, K. (1988). The structure of the human thyrotropin beta-subunit gene. *Gene* , 73 , 489-497.
11. Magner, J.A. (1990). Thyroid-stimulating hormone: biosynthesis, cell biology, and bioactivity. *Endocr Rev* , 11 , 354-385.
12. Magner, J.A. Biosynthesis, cell biology, and bioactivity of thyroid-stimulating hormone: update 1994. In: Braverman L, Refetoff S, editors. *Endocrine Reviews Monograph* . Rockville: The Endocrine Society, 1994:55.
13. Weintraub, B.D., Stannard, B.S., Magner, J.A., Ronin, C., Taylor, T., Joshi, L., Constant, R.B., Menezes-Ferreira, M.M., Petrick, P.& Gesundheit, N. (1985). Glycosylation and posttranslational processing of thyroid-stimulating hormone: clinical implications. *Recent Prog Horm Res* , 41 , 577-606.
14. Weintraub, B.D., Stannard, B.S., Linnekin, D.& Marshall, M. (1980). Relationship of glycosylation to de novo thyroid-stimulating hormone biosynthesis and secretion by mouse pituitary tumor cells. *J Biol Chem* , 255 , 5715-5723.
15. Szkudlinski, M.W., Teh, N.G., Grossmann, M., Tropea, J.E.& Weintraub, B.D. (1996). Engineering human glycoprotein hormone superactive analogues. *Nat Biotechnol* , 14 , 1257-1263.
16. Kleinau, G.& Krause, G. (2009). Thyrotropin and homologous glycoprotein hormone receptors: structural and functional aspects of extracellular signaling mechanisms. *Endocr Rev* , 30 , 133-151.
17. Amir, S.M., Kubota, K., Tramontano, D., Ingbar, S.H.& Keutmann, H.T. (1987). The carbohydrate moiety of bovine thyrotropin is essential for full bioactivity but not for receptor recognition. *Endocrinology* , 120 , 345-352.
18. Menezes-Ferreira, M.M., Petrick, P.A.& Weintraub, B.D. (1986). Regulation of thyrotropin (TSH) bioactivity by TSH-releasing hormone and thyroid hormone. *Endocrinology* , 118 , 2125-2130.
19. Taylor, T.& Weintraub, B.D. (1989). Altered thyrotropin (TSH) carbohydrate structures in hypothalamic hypothyroidism created by paraventricular nuclear lesions are corrected by in vivo TSH-releasing hormone administration. *Endocrinology* , 125 , 2198-2203.
20. Gesundheit, N., Fink, D.L., Silverman, L.A.& Weintraub, B.D. (1987). Effect of thyrotropin-releasing hormone on the carbohydrate structure of secreted mouse thyrotropin. Analysis by lectin affinity chromatography. *J Biol Chem* , 262 , 5197-5203.
21. Petersen, V.B., McGregor, A.M., Belchetz, P.E., Elkeles, R.S.& Hall, R. (1978). The secretion of thyrotrophin with impaired biological activity in patients with hypothalamic-pituitary disease. *Clin Endocrinol (Oxf)* , 8 , 397-402.
22. Faglia, G., Bitensky, L., Pinchera, A., Ferrari, C., Paracchi, A., Beck-Peccoz, P., Ambrosi, B.& Spada, A. (1979). Thyrotropin secretion in patients with central hypothyroidism: evidence for reduced

- biological activity of immunoreactive thyrotropin. *J Clin Endocrinol Metab* , 48 , 989-998.
23. Beck-Peccoz, P., Amr, S., Menezes-Ferreira, M.M., Faglia, G.& Weintraub, B.D. (1985). Decreased receptor binding of biologically inactive thyrotropin in central hypothyroidism. Effect of treatment with thyrotropin-releasing hormone. *N Engl J Med* , 312 , 1085-1090.
24. Beck-Peccoz, P.& Persani, L. (1994). Variable biological activity of thyroid-stimulating hormone. *Eur J Endocrinol* , 131 , 331-340.
25. Persani, L., Terzolo, M., Asteria, C., Orlandi, F., Angeli, A.& Beck-Peccoz, P. (1995). Circadian variations of thyrotropin bioactivity in normal subjects and patients with primary hypothyroidism. *J Clin Endocrinol Metab* , 80 , 2722-2728.
26. Persani, L. (1998). Hypothalamic thyrotropin-releasing hormone and thyrotropin biological activity. *Thyroid* , 8 , 941-946.
27. Hayashizaki, Y., Hiraoka, Y., Endo, Y., Miyai, K.& Matsubara, K. (1989). Thyroid-stimulating hormone (TSH) deficiency caused by a single base substitution in the CAGYC region of the beta-subunit. *Embo J* , 8 , 2291-2296.
28. Hayashizaki, Y., Hiraoka, Y., Tatsumi, K., Hashimoto, T., Furuyama, J., Miyai, K., Nishijo, K., Matsuura, M., Kohno, H., Labbe, A.& et al. (1990). Deoxyribonucleic acid analyses of five families with familial inherited thyroid stimulating hormone deficiency. *J Clin Endocrinol Metab* , 71 , 792-796.
29. Dacou-Voutetakis, C., Feltquate, D.M., Drakopoulou, M., Kourides, I.A.& Dracopoli, N.C. (1990). Familial hypothyroidism caused by a nonsense mutation in the thyroid-stimulating hormone beta-subunit gene. *Am J Hum Genet* , 46 , 988-993.
30. Medeiros-Neto, G., Herodotou, D.T., Rajan, S., Kommareddi, S., de Lacerda, L., Sandrini, R., Boguszewski, M.C., Hollenberg, A.N., Radovick, S.& Wondisford, F.E. (1996). A circulating, biologically inactive thyrotropin caused by a mutation in the beta subunit gene. *J Clin Invest* , 97 , 1250-1256.
31. Doeker, B.M., Pfaffle, R.W., Pohlenz, J.& Andler, W. (1998). Congenital central hypothyroidism due to a homozygous mutation in the thyrotropin beta-subunit gene follows an autosomal recessive inheritance. *J Clin Endocrinol Metab* , 83 , 1762-1765.
32. Heinrichs, C., Parma, J., Scherberg, N.H., Delange, F., Van Vliet, G., Duprez, L., Bourdoux, P., Bergmann, P., Vassart, G.& Refetoff, S. (2000). Congenital central isolated hypothyroidism caused by a homozygous mutation in the TSH-beta subunit gene. *Thyroid* , 10 , 387-391.
33. Bonomi, M., Proverbio, M.C., Weber, G., Chiumello, G., Beck-Peccoz, P.& Persani, L. (2001). Hyperplastic pituitary gland, high serum glycoprotein hormone alpha- subunit, and variable circulating thyrotropin (TSH) levels as hallmark of central hypothyroidism due to mutations of the TSH beta gene. *J Clin Endocrinol Metab* , 86 , 1600-1604.
34. Vallette-Kasic, S., Barlier, A., Teinturier, C., Diaz, A., Manavela, M., Berthezenne, F., Bouchard, P., Chaussain, J.L., Brauner, R., Pellegrini-Bouiller, I., Jaquet, P., Enjalbert, A.& Brue, T. (2001). PROP1 gene screening in patients with multiple pituitary hormone deficiency reveals two sites of hypermutability and a high incidence of corticotroph deficiency. *J Clin Endocrinol Metab* , 86 , 4529-4535.
35. Pohlenz, J., Dumitrescu, A., Aumann, U., Koch, G., Melchior, R., Prawitt, D.& Refetoff, S. (2002). Congenital secondary hypothyroidism caused by exon skipping due to a homozygous donor splice site mutation in the TSHbeta-subunit gene. *J Clin Endocrinol Metab* , 87 , 336-339.
36. McDermott, M.T., Haugen, B.R., Black, J.N., Wood, W.M., Gordon, D.F.& Ridgway, E.C. (2002).

Congenital isolated central hypothyroidism caused by a “hot spot” mutation in the thyrotropin-beta gene. *Thyroid* , 12 , 1141-1146.

37. Deladoey, J., Vuissoz, J.M., Domene, H.M., Malik, N., Gruneiro-Papendieck, L., Chiesa, A., Heinrich, J.J.& Mullis, P.E. (2003). Congenital secondary hypothyroidism due to a mutation C105Vfs114X thyrotropin-beta mutation: genetic study of five unrelated families from Switzerland and Argentina. *Thyroid* , 13 , 553-559.
38. Baquedano, M.S., Ciaccio, M., Dujovne, N., Herzovich, V., Longueira, Y., Warman, D.M., Rivarola, M.A.& Belgorosky, A. (2010). Two novel mutations of the TSH-beta subunit gene underlying congenital central hypothyroidism undetectable in neonatal TSH screening. *J Clin Endocrinol Metab* , 95 , E98-103.
39. Azzam, N., Bar-Shalom, R., Kraiem, Z.& Fares, F. (2005). Human thyrotropin (TSH) variants designed by site-directed mutagenesis block TSH activity in vitro and in vivo. *Endocrinology* , 146 , 2845-2850.
40. Mueller, S., Kleinau, G., Szkudlinski, M.W., Jaeschke, H., Krause, G.& Paschke, R. (2009). The superagonistic activity of bovine thyroid-stimulating hormone (TSH) and the human TR1401 TSH analog is determined by specific amino acids in the hinge region of the human TSH receptor. *J Biol Chem* , 284 , 16317-16324.
41. Nakabayashi, K., Matsumi, H., Bhalla, A., Bae, J., Mosselman, S., Hsu, S.Y.& Hsueh, A.J. (2002). Thyrostimulin, a heterodimer of two new human glycoprotein hormone subunits, activates the thyroid-stimulating hormone receptor. *J Clin Invest* , 109 , 1445-1452.
42. Nagasaki, H., Wang, Z., Jackson, V.R., Lin, S., Nothacker, H.P.& Civelli, O. (2006). Differential expression of the thyrostimulin subunits, glycoprotein alpha2 and beta5 in the rat pituitary. *J Mol Endocrinol* , 37 , 39-50.
43. Wondisford, F.E. (2002). The thyroid axis just got more complicated. *J Clin Invest* , 109 , 1401-1402.
44. Sun, S.C., Hsu, P.J., Wu, F.J., Li, S.H., Lu, C.H.& Luo, C.W. (2010). Thyrostimulin, but not thyroid-stimulating hormone (TSH), acts as a paracrine regulator to activate the TSH receptor in mammalian ovary. *J Biol Chem* , 285 , 3758-3765.
45. Reichlin, S.& Utiger, R.D. (1967). Regulation of the pituitary-thyroid axis in man: relationship of TSH concentration to concentration of free and total thyroxine in plasma. *J Clin Endocrinol Metab* , 27 , 251-255.
46. Bogazzi, F., Bartalena, L., Brogioni, S., Burelli, A., Grasso, L., Dell'Unto, E., Manetti, L.& Martino, E. (1997). L-thyroxine directly affects expression of thyroid hormone-sensitive genes: regulatory effect of RXRbeta. *Mol Cell Endocrinol* , 134 , 23-31.
47. Gurr, J.A.& Kourides, I.A. (1983). Regulation of thyrotropin biosynthesis. Discordant effect of thyroid hormone on alpha and beta subunit mRNA levels. *J Biol Chem* , 258 , 10208-10211.
48. Shupnik, M.A., Chin, W.W., Habener, J.F.& Ridgway, E.C. (1985). Transcriptional regulation of the thyrotropin subunit genes by thyroid hormone. *J Biol Chem* , 260 , 2900-2903.
49. Ridgway, E.C., Kourides, I.A., Chin, W.W., Cooper, D.S.& Maloof, F. (1979). Augmentation of pituitary thyrotrophin response to thyrotrophin releasing hormone during subphysiological tri-iodothyroinone therapy in hypothyroidism. *Clin Endocrinol (Oxf)* , 10 , 343-353.
50. Chin, W.W., Carr, F.E., Burnside, J.& Darling, D.S. (1993). Thyroid hormone regulation of thyrotropin gene expression. *Recent Prog Horm Res* , 48 , 393-414.

51. Shupnik, M.A.& Ridgway, E.C. (1987). Thyroid hormone control of thyrotropin gene expression in rat anterior pituitary cells. *Endocrinology* , 121 , 619-624.
52. Kourides, I.A., Gurr, J.A.& Wolf, O. (1984). The regulation and organization of thyroid stimulating hormone genes. *Recent Prog Horm Res* , 40 , 79-120.
53. Carr, F.E., Shupnik, M.A., Burnside, J.& Chin, W.W. (1989). Thyrotropin-releasing hormone stimulates the activity of the rat thyrotropin beta-subunit gene promoter transfected into pituitary cells. *Mol Endocrinol* , 3 , 717-724.
54. Carr, F.E., Ridgway, E.C.& Chin, W.W. (1985). Rapid simultaneous measurement of rat alpha- and thyrotropin (TSH) beta-subunit messenger ribonucleic acids (mRNAs) by solution hybridization: regulation of TSH subunit mRNAs by thyroid hormones. *Endocrinology* , 117 , 1272-1278.
55. Gurr, J.A., Januszkesi, M.M., Tidikis, I.M., Norcross, J.J.& Kourides, I.A. (1990). Thyroid hormone regulates expression of the thyrotropin beta-subunit gene from both transcription start sites in the mouse and rat. *Mol Cell Endocrinol* , 71 , 185-193.
56. Chatterjee, V.K., Lee, J.K., Rentoumis, A.& Jameson, J.L. (1989). Negative regulation of the thyroid-stimulating hormone alpha gene by thyroid hormone: receptor interaction adjacent to the TATA box. *Proc Natl Acad Sci U S A* , 86 , 9114-9118.
57. Wondisford, F.E., Farr, E.A., Radovick, S., Steinfeldler, H.J., Moates, J.M., McClaskey, J.H.& Weintraub, B.D. (1989). Thyroid hormone inhibition of human thyrotropin beta-subunit gene expression is mediated by a cis-acting element located in the first exon. *J Biol Chem* , 264 , 14601-14604.
58. Bodenner, D.L., Mroczynski, M.A., Weintraub, B.D., Radovick, S.& Wondisford, F.E. (1991). A detailed functional and structural analysis of a major thyroid hormone inhibitory element in the human thyrotropin beta-subunit gene. *J Biol Chem* , 266 , 21666-21673.
59. Darling, D.S., Burnside, J.& Chin, W.W. (1989). Binding of thyroid hormone receptors to the rat thyrotropin-beta gene. *Mol Endocrinol* , 3 , 1359-1368.
60. Brent, G.A., Williams, G.R., Harney, J.W., Forman, B.M., Samuels, H.H., Moore, D.D.& Larsen, P.R. (1991). Effects of varying the position of thyroid hormone response elements within the rat growth hormone promoter: implications for positive and negative regulation by 3,5,3'-triiodothyronine. *Mol Endocrinol* , 5 , 542-548.
61. Carr, F.E.& Wong, N.C. (1994). Characteristics of a negative thyroid hormone response element. *J Biol Chem* , 269 , 4175-4179.
62. Burnside, J., Darling, D.S., Carr, F.E.& Chin, W.W. (1989). Thyroid hormone regulation of the rat glycoprotein hormone alpha-subunit gene promoter activity. *J Biol Chem* , 264 , 6886-6891.
63. Shupnik, M.A., Greenspan, S.L.& Ridgway, E.C. (1986). Transcriptional regulation of thyrotropin subunit genes by thyrotropin-releasing hormone and dopamine in pituitary cell culture. *J Biol Chem* , 261 , 12675-12679.
64. Langlois, M.F., Zanger, K., Monden, T., Safer, J.D., Hollenberg, A.N.& Wondisford, F.E. (1997). A unique role of the beta-2 thyroid hormone receptor isoform in negative regulation by thyroid hormone. Mapping of a novel amino-terminal domain important for ligand-independent activation. *J Biol Chem* , 272 , 24927-24933.
65. Wikstrom, L., Johansson, C., Salto, C., Barlow, C., Campos Barros, A., Baas, F., Forrest, D., Thoren, P.& Vennstrom, B. (1998). Abnormal heart rate and body temperature in mice lacking thyroid hormone receptor alpha 1. *Embo J* , 17 , 455-461.

66. Forrest, D., Hanebuth, E., Smeyne, R.J., Everds, N., Stewart, C.L., Wehner, J.M. & Curran, T. (1996). Recessive resistance to thyroid hormone in mice lacking thyroid hormone receptor beta: evidence for tissue-specific modulation of receptor function. *Embo J* , 15 , 3006-3015.
67. Weiss, R.E., Forrest, D., Pohlenz, J., Cua, K., Curran, T. & Refetoff, S. (1997). Thyrotropin regulation by thyroid hormone in thyroid hormone receptor beta-deficient mice. *Endocrinology* , 138 , 3624-3629.
68. Abel, E.D., Boers, M.E., Pazos-Moura, C., Moura, E., Kaulbach, H., Zakaria, M., Lowell, B., Radovick, S., Liberman, M.C. & Wondisford, F. (1999). Divergent roles for thyroid hormone receptor beta isoforms in the endocrine axis and auditory system. *J Clin Invest* , 104 , 291-300.
69. Gothe, S., Wang, Z., Ng, L., Kindblom, J.M., Barros, A.C., Ohlsson, C., Vennstrom, B. & Forrest, D. (1999). Mice devoid of all known thyroid hormone receptors are viable but exhibit disorders of the pituitary-thyroid axis, growth, and bone maturation. *Genes Dev* , 13 , 1329-1341.
70. Gauthier, K., Chassande, O., Plateroti, M., Roux, J.P., Legrand, C., Pain, B., Roussel, B., Weiss, R., Trouillas, J. & Samarut, J. (1999). Different functions for the thyroid hormone receptors TRalpha and TRbeta in the control of thyroid hormone production and post-natal development. *Embo J* , 18 , 623-631.
71. Hayashi, Y., Xie, J., Weiss, R.E., Pohlenz, J. & Refetoff, S. (1998). Selective pituitary resistance to thyroid hormone produced by expression of a mutant thyroid hormone receptor beta gene in the pituitary gland of transgenic mice. *Biochem Biophys Res Commun* , 245 , 204-210.
72. Weiss, R.E., Chassande, O., Koo, E.K., Macchia, P.E., Cua, K., Samarut, J. & Refetoff, S. (2002). Thyroid function and effect of aging in combined hetero/homozygous mice deficient in thyroid hormone receptors alpha and beta genes. *J Endocrinol* , 172 , 177-185.
73. Abel, E.D., Moura, E.G., Ahima, R.S., Campos-Barros, A., Pazos-Moura, C.C., Boers, M.E., Kaulbach, H.C., Forrest, D. & Wondisford, F.E. (2003). Dominant inhibition of thyroid hormone action selectively in the pituitary of thyroid hormone receptor-beta null mice abolishes the regulation of thyrotropin by thyroid hormone. *Mol Endocrinol* , 17 , 1767-1776.
74. Zenke, M., Munoz, A., Sap, J., Vennstrom, B. & Beug, H. (1990). v-erbA oncogene activation entails the loss of hormone-dependent regulator activity of c-erbA. *Cell* , 61 , 1035-1049.
75. Morita, S., Fernandez-Mejia, C. & Melmed, S. (1989). Retinoic acid selectively stimulates growth hormone secretion and messenger ribonucleic acid levels in rat pituitary cells. *Endocrinology* , 124 , 2052-2056.
76. Wondisford, F.E., Steinfelder, H.J., Nations, M. & Radovick, S. (1993). AP-1 antagonizes thyroid hormone receptor action on the thyrotropin beta-subunit gene. *J Biol Chem* , 268 , 2749-2754.
77. Shibusawa, N., Hollenberg, A.N. & Wondisford, F.E. (2003). Thyroid hormone receptor DNA binding is required for both positive and negative gene regulation. *J Biol Chem* , 278 , 732-738.
78. Nakano, K., Matsushita, A., Sasaki, S., Misawa, H., Nishiyama, K., Kashiwabara, Y. & Nakamura, H. (2004). Thyroid-hormone-dependent negative regulation of thyrotropin beta gene by thyroid hormone receptors: study with a new experimental system using CV1 cells. *Biochem J* , 378 , 549-557.
79. Hallenbeck, P.L., Phyllaier, M. & Nikodem, V.M. (1993). Divergent effects of 9-cis-retinoic acid receptor on positive and negative thyroid hormone receptor-dependent gene expression. *J Biol Chem* , 268 , 3825-3828.
80. Cohen, O., Flynn, T.R. & Wondisford, F.E. (1995). Ligand-dependent antagonism by retinoid X receptors of inhibitory thyroid hormone response elements. *J Biol Chem* , 270 , 13899-13905.

81. Breen, J.J., Hickok, N.J.& Gurr, J.A. (1997). The rat TSHbeta gene contains distinct response elements for regulation by retinoids and thyroid hormone. *Mol Cell Endocrinol* , 131 , 137-146.
82. Haugen, B.R., Brown, N.S., Wood, W.M., Gordon, D.F.& Ridgway, E.C. (1997). The thyrotrope-restricted isoform of the retinoid-X receptor-gamma1 mediates 9-cis-retinoic acid suppression of thyrotropin-beta promoter activity. *Mol Endocrinol* , 11 , 481-489.
83. Cheng, S.Y., Leonard, J.L.& Davis, P.J. (2010). Molecular aspects of thyroid hormone actions. *Endocr Rev* , 31 , 139-170.
84. Kim, M.K., McClaskey, J.H., Bodenner, D.L.& Weintraub, B.D. (1993). A c-fos/c-jun-like factor and the pituitary-specific factor Pit-1 are both necessary to mediate induction of the human thyrotropin beta promoter. *Trans Assoc Am Physicians* , 106 , 62-68.
85. Pfahl, M. (1993). Nuclear receptor/AP-1 interaction. *Endocr Rev* , 14 , 651-658.
86. Koenig, R.J.& Watson, A.Y. (1984). Enrichment of rat anterior pituitary cell types by metrizamide density gradient centrifugation. *Endocrinology* , 115 , 317-323.
87. Farquhar, R.& Rinehart, J. (1954). Cytological alterations in the anterior pituitary gland following thyroideectomy. *Endocrinology* , 55 , 857.
88. Larsen, P.R., Silva, J.E.& Kaplan, M.M. (1981). Relationships between circulating and intracellular thyroid hormones: physiological and clinical implications. *Endocr Rev* , 2 , 87-102.
89. Chin, W.W., Shupnik, M.A., Ross, D.S., Habener, J.F.& Ridgway, E.C. (1985). Regulation of the alpha and thyrotropin beta-subunit messenger ribonucleic acids by thyroid hormones. *Endocrinology* , 116 , 873-878.
90. Franklyn, J.A., Wood, D.F., Balfour, N.J., Ramsden, D.B., Docherty, K., Chin, W.W.& Sheppard, M.C. (1987). Effect of hypothyroidism and thyroid hormone replacement in vivo on pituitary cytoplasmic concentrations of thyrotropin-beta and alpha-subunit messenger ribonucleic acids. *Endocrinology* , 120 , 2279-2288.
91. Silva, J.E.& Larsen, P.R. (1977). Pituitary nuclear 3,5,3'-triiodothyronine and thyrotropin secretion: an explanation for the effect of thyroxine. *Science* , 198 , 617-620.
92. Vale, W., Burgus, R.& Guillemin, R. (1968). On the mechanism of action of TRF: effects of cycloheximide and actinomycin on the release of TSH stimulated in vitro by TRF and its inhibition by thyroxine. *Neuroendocrinology* , 3 , 34.
93. Bowers, C.Y., Lee, K.L.& Schally, A.V. (1968). A study on the interaction of the thyrotropin-releasing factor and L-triiodothyronine: effects of puromycin and cycloheximide. *Endocrinology* , 82 , 75-82.
94. Gard, T.G., Bernstein, B.& Larsen, P.R. (1981). Studies on the mechanism of 3,5,3'-triiodothyronine-induced suppression of secretagogue-induced thyrotropin release in vitro. *Endocrinology* , 108 , 2046-2053.
95. Koenig, R.J., Senator, D.& Larsen, P.R. (1984). Phorbol esters as probes of the regulation of thyrotropin secretion. *Biochem Biophys Res Commun* , 125 , 353-359.
96. Schrey, M.P.& Larsen, P.R. (1981). Evidence for a possible role for Ca++ in the 3,5,3'-triiodothyronine inhibition of thyrotropin-releasing hormone-induced secretion of thyrotropin by rat anterior pituitary in vitro. *Endocrinology* , 108 , 1690-1696.
97. Obregon, M.J., Pascual, A., Mallol, J., Morreale de Escobar, G.& Escobar del Rey, F. (1980). Evidence against a major role of L-thyroxine at the pituitary level: studies in rats treated with iopanoic

- acid (telepaque). *Endocrinology* , 106 , 1827-1836.
98. Lueprasitsakul, W., Alex, S., Fang, S.L., Pino, S., Irmscher, K., Kohrle, J.& Braverman, L.E. (1990). Flavonoid administration immediately displaces thyroxine (T4) from serum transthyretin, increases serum free T4, and decreases serum thyrotropin in the rat. *Endocrinology* , 126 , 2890-2895.
99. Spencer, C.A., Takeuchi, M., Kazarosyan, M., MacKenzie, F., Beckett, G.J.& Wilkinson, E. (1995). Interlaboratory/intermethod differences in functional sensitivity of immunometric assays of thyrotropin (TSH) and impact on reliability of measurement of subnormal concentrations of TSH. *Clin Chem* , 41 , 367-374.
100. Riesco, G., Taurog, A., Larsen, R.& Krulich, L. (1977). Acute and chronic responses to iodine deficiency in rats. *Endocrinology* , 100 , 303-313.
101. Patel, Y.C., Pharoah, P.O., Hornabrook, R.W.& Hetzel, B.S. (1973). Serum triiodothyronine, thyroxine and thyroid-stimulating hormone in endemic goiter: a comparison of goitrous and nongoitrous subjects in New Guinea. *J Clin Endocrinol Metab* , 37 , 783-789.
102. Wennlund, A. (1986). Variation in serum levels of T3, T4, FT4 and TSH during thyroxine replacement therapy. *Acta Endocrinol (Copenh)* , 113 , 47-49.
103. Bigos, S.T., Ridgway, E.C., Kourides, I.A.& Maloof, F. (1978). Spectrum of pituitary alterations with mild and severe thyroid impairment. *J Clin Endocrinol Metab* , 46 , 317-325.
104. Jackson, I.M. (1982). Thyrotropin-releasing hormone. *N Engl J Med* , 306 , 145-155.
105. Guillemin, R. (2005). Hypothalamic hormones a.k.a. hypothalamic releasing factors. *J Endocrinol* , 184 , 11-28.
106. Jeffcoate, S.L., White, N., Hokfelt, T., Fuxe, K.& Johansson, O. (1976). Proceedings: Localization of thyrotrophin releasing hormone in the spinal cord of the rat by immunohisto-chemistry and radioimmunoassay. *J Endocrinol* , 69 , 9P-10P.
107. Martino, E., Lernmark, A., Seo, H., Steiner, D.F.& Refetoff, S. (1978). High concentration of thyrotropin-releasing hormone in pancreatic islets. *Proc Natl Acad Sci U S A* , 75 , 4265-4267.
108. Gkonos, P.J., Tavianini, M.A., Liu, C.C.& Roos, B.A. (1989). Thyrotropin-releasing hormone gene expression in normal thyroid parafollicular cells. *Mol Endocrinol* , 3 , 2101-2109.
109. Bruhn, T.O., Rondeel, J.M., Bolduc, T.G.& Jackson, I.M. (1994). Thyrotropin-releasing hormone (TRH) gene expression in the anterior pituitary. I. Presence of pro-TRH messenger ribonucleic acid and pro-TRH-derived peptide in a subpopulation of somatotrophs. *Endocrinology* , 134 , 815-820.
110. Bilek, R. (2000). TRH-like peptides in prostate gland and other tissues. *Physiol Res* , 49 , S19-26.
111. Bruhn, T.O., Huang, S.S., Vaslet, C.& Nillni, E.A. (1998). Glucocorticoids modulate the biosynthesis and processing of prothyrotropin releasing-hormone (proTRH). *Endocrine* , 9 , 143-152.
112. Fekete, C.& Lechan, R.M. (2007). Negative feedback regulation of hypophysiotropic thyrotropin-releasing hormone (TRH) synthesizing neurons: role of neuronal afferents and type 2 deiodinase. *Front Neuroendocrinol* , 28 , 97-114.
113. Bruhn, T.O., Rondeel, J.M.& Jackson, I.M. (1998). Thyrotropin-releasing hormone gene expression in the anterior pituitary. IV. Evidence for paracrine and autocrine regulation. *Endocrinology* , 139 , 3416-3422.
114. Lee, S.L., Stewart, K.& Goodman, R.H. (1988). Structure of the gene encoding rat thyrotropin releasing hormone. *J Biol Chem* , 263 , 16604-16609.

115. Yamada, M., Radovick, S., Wondisford, F.E., Nakayama, Y., Weintraub, B.D.& Wilber, J.F. (1990). Cloning and structure of human genomic DNA and hypothalamic cDNA encoding human prepro thyrotropin-releasing hormone. *Mol Endocrinol* , 4 , 551-556.
116. Yamada, M., Wondisford, F.E., Radovick, S., Nakayama, Y., Weintraub, B.D.& Wilber, J.F. (1991). Assignment of human preprothyrotropin-releasing hormone (TRH) gene to chromosome 3. *Somat Cell Mol Genet* , 17 , 97-100.
117. Bulant, M., Roussel, J.P., Astier, H., Nicolas, P.& Vaudry, H. (1990). Processing of thyrotropin-releasing hormone prohormone (pro-TRH) generates a biologically active peptide, prepro-TRH-(160-169), which regulates TRH-induced thyrotropin secretion. *Proc Natl Acad Sci U S A* , 87 , 4439-4443.
118. Pekary, A.E. (1998). Is Ps4 (prepro-TRH [160-169]) more than an enhancer for thyrotropin-releasing hormone? *Thyroid* , 8 , 963-968.
119. Nillni, E.A. (1999). Neuroregulation of ProTRH biosynthesis and processing. *Endocrine* , 10 , 185-199.
120. Carr, F.E., Fein, H.G., Fisher, C.U., Wessendorf, M.W.& Smallridge, R.C. (1992). A cryptic peptide (160-169) of thyrotropin-releasing hormone prohormone demonstrates biological activity in vivo and in vitro. *Endocrinology* , 131 , 2653-2658.
121. Carr, F.E., Reid, A.H.& Wessendorf, M.W. (1993). A cryptic peptide from the preprothyrotropin-releasing hormone precursor stimulates thyrotropin gene expression. *Endocrinology* , 133 , 809-814.
122. Engler, D., Redei, E.& Kola, I. (1999). The corticotropin-release inhibitory factor hypothesis: a review of the evidence for the existence of inhibitory as well as stimulatory hypophysiotropic regulation of adrenocorticotropin secretion and biosynthesis. *Endocr Rev* , 20 , 460-500.
123. Jessop, D.S. (1999). Review: Central non-glucocorticoid inhibitors of the hypothalamo-pituitary-adrenal axis. *J Endocrinol* , 160 , 169-180.
124. Nillni, E.A.& Sevarino, K.A. (1999). The biology of pro-thyrotropin-releasing hormone-derived peptides. *Endocr Rev* , 20 , 599-648.
125. Segerson, T.P., Kauer, J., Wolfe, H.C., Mobtaker, H., Wu, P., Jackson, I.M.& Lechan, R.M. (1987). Thyroid hormone regulates TRH biosynthesis in the paraventricular nucleus of the rat hypothalamus. *Science* , 238 , 78-80.
126. Koller, K.J., Wolff, R.S., Warden, M.K.& Zoeller, R.T. (1987). Thyroid hormones regulate levels of thyrotropin-releasing-hormone mRNA in the paraventricular nucleus. *Proc Natl Acad Sci U S A* , 84 , 7329-7333.
127. Dyess, E.M., Segerson, T.P., Liposits, Z., Paull, W.K., Kaplan, M.M., Wu, P., Jackson, I.M.& Lechan, R.M. (1988). Triiodothyronine exerts direct cell-specific regulation of thyrotropin-releasing hormone gene expression in the hypothalamic paraventricular nucleus. *Endocrinology* , 123 , 2291-2297.
128. Kakucska, I., Rand, W.& Lechan, R.M. (1992). Thyrotropin-releasing hormone gene expression in the hypothalamic paraventricular nucleus is dependent upon feedback regulation by both triiodothyronine and thyroxine. *Endocrinology* , 130 , 2845-2850.
129. Strait, K.A., Zou, L.& Oppenheimer, J.H. (1992). Beta 1 isoform-specific regulation of a triiodothyronine-induced gene during cerebellar development. *Mol Endocrinol* , 6 , 1874-1880.
130. Lechan, R.M., Qi, Y., Jackson, I.M.& Mahdavi, V. (1994). Identification of thyroid hormone receptor isoforms in thyrotropin-releasing hormone neurons of the hypothalamic paraventricular

- nucleus. *Endocrinology* , 135 , 92-100.
131. Wilber, J.F.& Xu, A.H. (1998). The thyrotropin-releasing hormone gene 1998: cloning, characterization, and transcriptional regulation in the central nervous system, heart, and testis. *Thyroid* , 8 , 897-901.
132. Abel, E.D., Ahima, R.S., Boers, M.E., Elmquist, J.K.& Wondisford, F.E. (2001). Critical role for thyroid hormone receptor beta2 in the regulation of paraventricular thyrotropin-releasing hormone neurons. *J Clin Invest* , 107 , 1017-1023.
133. Riskind, P.N., Kolodny, J.M.& Larsen, P.R. (1987). The regional hypothalamic distribution of type II 5'-monodeiodinase in euthyroid and hypothyroid rats. *Brain Res* , 420 , 194-198.
134. Rodriguez, E.M., Blazquez, J.L., Pastor, F.E., Pelaez, B., Pena, P., Peruzzo, B.& Amat, P. (2005). Hypothalamic tanycytes: a key component of brain-endocrine interaction. *Int Rev Cytol* , 247 , 89-164.
135. Jansen, J., Friesema, E.C., Milici, C.& Visser, T.J. (2005). Thyroid hormone transporters in health and disease. *Thyroid* , 15 , 757-768.
136. Yamada, M., Satoh, T., Monden, T.& Mori, M. (1999). Assignment of the thyrotropin-releasing hormone gene (TRH) to human chromosome 3q13.3->q21 by in situ hybridization. *Cytogenet Cell Genet* , 87 , 275.
137. Chiamolera, M.I.& Wondisford, F.E. (2009). Minireview: Thyrotropin-releasing hormone and the thyroid hormone feedback mechanism. *Endocrinology* , 150 , 1091-1096.
138. Nikrothananond, A.A., Ortiga-Carvalho, T.M., Shibusawa, N., Hashimoto, K., Liao, X.H., Refetoff, S., Yamada, M., Mori, M.& Wondisford, F.E. (2006). Dominant role of thyrotropin-releasing hormone in the hypothalamic-pituitary-thyroid axis. *J Biol Chem* , 281 , 5000-5007.
139. Mittag, J., Friedrichsen, S., Strube, A., Heuer, H.& Bauer, K. (2009). Analysis of hypertrophic thyrotrophs in pituitaries of athyroid Pax8-/- mice. *Endocrinology* , 150 , 4443-4449.
140. Cintra, A., Fuxe, K., Wikstrom, A.C., Visser, T.& Gustafsson, J.A. (1990). Evidence for thyrotropin-releasing hormone and glucocorticoid receptor-immunoreactive neurons in various preoptic and hypothalamic nuclei of the male rat. *Brain Res* , 506 , 139-144.
141. Jackson, I.M. (1995). Thyrotropin-releasing hormone and corticotropin-releasing hormone—what's the message? *Endocrinology* , 136 , 2793-2794.
142. Luo, L.G., Bruhn, T.& Jackson, I.M. (1995). Glucocorticoids stimulate thyrotropin-releasing hormone gene expression in cultured hypothalamic neurons. *Endocrinology* , 136 , 4945-4950.
143. Shibusawa, N., Yamada, M., Hirato, J., Monden, T., Satoh, T.& Mori, M. (2000). Requirement of thyrotropin-releasing hormone for the postnatal functions of pituitary thyrotrophs: ontogeny study of congenital tertiary hypothyroidism in mice. *Mol Endocrinol* , 14 , 137-146.
144. Straub, R.E., Frech, G.C., Joho, R.H.& Gershengorn, M.C. (1990). Expression cloning of a cDNA encoding the mouse pituitary thyrotropin-releasing hormone receptor. *Proc Natl Acad Sci U S A* , 87 , 9514-9518.
145. Hinuma, S., Hosoya, M., Ogi, K., Tanaka, H., Nagai, Y.& Onda, H. (1994). Molecular cloning and functional expression of a human thyrotropin-releasing hormone (TRH) receptor gene. *Biochim Biophys Acta* , 1219 , 251-259.
146. Bilek, R.& Starka, L. (2005). The computer modelling of human TRH receptor, TRH and TRH-like peptides. *Physiol Res* , 54 , 141-150.

147. Collu, R., Tang, J., Castagne, J., Lagace, G., Masson, N., Huot, C., Deal, C., Delvin, E., Faccenda, E., Eidne, K.A.& Van Vliet, G. (1997). A novel mechanism for isolated central hypothyroidism: inactivating mutations in the thyrotropin-releasing hormone receptor gene. *J Clin Endocrinol Metab* , 82 , 1561-1565.
148. Collu, R. (2000). Genetic aspects of central hypothyroidism. *J Endocrinol Invest* , 23 , 125-134.
149. Gershengorn, M.C. (1978). Bihormonal regulation of the thyrotropin-releasing hormone receptor in mouse pituitary thyrotropic tumor cells in culture. *J Clin Invest* , 62 , 937-943.
150. Fujimoto, J.& Gershengorn, M.C. (1991). Evidence for dual regulation by protein kinases A and C of thyrotropin-releasing hormone receptor mRNA in GH3 cells. *Endocrinology* , 129 , 3430-3432.
151. Brenner-Gati, L.& Gershengorn, M.C. (1986). Effects of thyrotropin-releasing hormone on phosphoinositides and cytoplasmic free calcium in thyrotropic pituitary cells. *Endocrinology* , 118 , 163-169.
152. Oron, Y., Gillo, B., Straub, R.E.& Gershengorn, M.C. (1987). Mechanism of membrane electrical response to thyrotropin-releasing hormone in Xenopus oocytes injected with GH3 pituitary cell messenger ribonucleic acid. *Mol Endocrinol* , 1 , 918-925.
153. Gershengorn, M.C. (1985). Thyrotropin-releasing hormone action: mechanism of calcium-mediated stimulation of prolactin secretion. *Recent Prog Horm Res* , 41 , 607-653.
154. Heinflink, M., Nussenzeig, D.R., Friedman, A.M.& Gershengorn, M.C. (1994). Thyrotropin-releasing hormone receptor activation does not elevate intracellular cyclic adenosine 3',5'-monophosphate in cells expressing high levels of receptors. *J Clin Endocrinol Metab* , 79 , 650-652.
155. Martin, T.F. (1983). Thyrotropin-releasing hormone rapidly activates the phosphodiester hydrolysis of polyphosphoinositides in GH3 pituitary cells. Evidence for the role of a polyphosphoinositide-specific phospholipase C in hormone action. *J Biol Chem* , 258 , 14816-14822.
156. Nelson, E.J.& Hinkle, P.M. (1994). Characteristics of the Ca<sup>2+</sup> spike and oscillations induced by different doses of thyrotropin-releasing hormone (TRH) in individual pituitary cells and nonexcitable cells transfected with TRH receptor complementary deoxyribonucleic acid. *Endocrinology* , 135 , 1084-1092.
157. Nelson, E.J.& Hinkle, P.M. (1994). Thyrotropin-releasing hormone activates Ca<sup>2+</sup> efflux. Evidence suggesting that a plasma membrane Ca<sup>2+</sup> pump is an effector for a G-protein-coupled Ca(2+)-mobilizing receptor. *J Biol Chem* , 269 , 30854-30860.
158. Ashworth, R.& Hinkle, P.M. (1996). Thyrotropin-releasing hormone-induced intracellular calcium responses in individual rat lactotrophs and thyrotrophs. *Endocrinology* , 137 , 5205-5212.
159. Carr, F.E., Burnside, J.& Chin, W.W. (1989). Thyroid hormones regulate rat thyrotropin beta gene promoter activity expressed in GH3 cells. *Mol Endocrinol* , 3 , 709-716.
160. Shupnik, M.A., Weck, J.& Hinkle, P.M. (1996). Thyrotropin (TSH)-releasing hormone stimulates TSH beta promoter activity by two distinct mechanisms involving calcium influx through L type Ca<sup>2+</sup> channels and protein kinase C. *Mol Endocrinol* , 10 , 90-99.
161. Steinfelder, H.J.& Wondisford, F.E. (1997). Thyrotropin (TSH) beta-subunit gene expression—an example for the complex regulation of pituitary hormone genes. *Exp Clin Endocrinol Diabetes* , 105 , 196-203.
162. Steinfelder, H.J., Hauser, P., Nakayama, Y., Radovick, S., McClaskey, J.H., Taylor, T., Weintraub, B.D.& Wondisford, F.E. (1991). Thyrotropin-releasing hormone regulation of human TSHB

- expression: role of a pituitary-specific transcription factor (Pit-1/GHF-1) and potential interaction with a thyroid hormone-inhibitory element. *Proc Natl Acad Sci U S A* , 88 , 3130-3134.
163. Weintraub, B.D., Wondisford, F.E., Farr, E.A., Steinfeldler, H.J., Radovick, S., Gesundheit, N., Gyves, P.W., Taylor, T.& DeCherney, G.S. (1989). Pre-translational and post-translational regulation of TSH synthesis in normal and neoplastic thyrotrophs. *Horm Res* , 32 , 22-24.
164. Heuer, H., Schafer, M.K.& Bauer, K. (1998). The thyrotropin-releasing hormone-degrading ectoenzyme: the third element of the thyrotropin-releasing hormone-signaling system. *Thyroid* , 8 , 915-920.
165. Steinfeldler, H.J., Radovick, S., Mroczynski, M.A., Hauser, P., McClaskey, J.H., Weintraub, B.D.& Wondisford, F.E. (1992). Role of a pituitary-specific transcription factor (pit-1/GHF-1) or a closely related protein in cAMP regulation of human thyrotropin-beta subunit gene expression. *J Clin Invest* , 89 , 409-419.
166. Steinfeldler, H.J., Radovick, S.& Wondisford, F.E. (1992). Hormonal regulation of the thyrotropin beta-subunit gene by phosphorylation of the pituitary-specific transcription factor Pit-1. *Proc Natl Acad Sci U S A* , 89 , 5942-5945.
167. Howard, P.W.& Maurer, R.A. (1994). Thyrotropin releasing hormone stimulates transient phosphorylation of the tissue-specific transcription factor, Pit-1. *J Biol Chem* , 269 , 28662-28669.
168. Parks, J.S., Brown, M.R., Hurley, D.L., Phelps, C.J.& Wajnrajch, M.P. (1999). Heritable disorders of pituitary development. *J Clin Endocrinol Metab* , 84 , 4362-4370.
169. Camper, S.A., Saunders, T.L., Katz, R.W.& Reeves, R.H. (1990). The Pit-1 transcription factor gene is a candidate for the murine Snell dwarf mutation. *Genomics* , 8 , 586-590.
170. Li, S., Crenshaw, E.B., 3rd, Rawson, E.J., Simmons, D.M., Swanson, L.W.& Rosenfeld, M.G. (1990). Dwarf locus mutants lacking three pituitary cell types result from mutations in the POU-domain gene pit-1. *Nature* , 347 , 528-533.
171. Parks, J.S., Kinoshita, E.I.& Pfaffle, R.W. (1993). Pit-1 and hypopituitarism. *TEM* , 4 , 81.
172. Radovick, S., Nations, M., Du, Y., Berg, L.A., Weintraub, B.D.& Wondisford, F.E. (1992). A mutation in the POU-homeodomain of Pit-1 responsible for combined pituitary hormone deficiency. *Science* , 257 , 1115-1118.
173. Pfaffle, R.W., DiMatta, G.E., Parks, J.S., Brown, M.R., Wit, J.M., Jansen, M., Van der Nat, H., Van den Brande, J.L., Rosenfeld, M.G.& Ingraham, H.A. (1992). Mutation of the POU-specific domain of Pit-1 and hypopituitarism without pituitary hypoplasia. *Science* , 257 , 1118-1121.
174. Tatsumi, K., Miyai, K., Notomi, T., Kaibe, K., Amino, N., Mizuno, Y.& Kohno, H. (1992). Cretinism with combined hormone deficiency caused by a mutation in the PIT1 gene. *Nat Genet* , 1 , 56-58.
175. Turton, J.P., Reynaud, R., Mehta, A., Torpiano, J., Saveanu, A., Woods, K.S., Tiulpakov, A., Zdravkovic, V., Hamilton, J., Attard-Montalto, S., Parascandalo, R., Vella, C., Clayton, P.E., Shalet, S., Barton, J., Brue, T.& Dattani, M.T. (2005). Novel mutations within the POU1F1 gene associated with variable combined pituitary hormone deficiency. *J Clin Endocrinol Metab* , 90 , 4762-4770.
176. Hashimoto, Y., Cisternino, M.& Cohen, L.E. (2003). A novel nonsense mutation in the Pit-1 gene: evidence for a gene dosage effect. *J Clin Endocrinol Metab* , 88 , 1241-1247.
177. Cohen, L.E., Wondisford, F.E.& Radovick, S. (1996). Role of Pit-1 in the gene expression of growth hormone, prolactin, and thyrotropin. *Endocrinol Metab Clin North Am* , 25 , 523-540.

178. Gordon, D.F., Haugen, B.R., Sarapura, V.D., Nelson, A.R., Wood, W.M.& Ridgway, E.C. (1993). Analysis of Pit-1 in regulating mouse TSH beta promoter activity in thyrotropes. *Mol Cell Endocrinol* , 96 , 75-84.
179. Haugen, B.R., Wood, W.M., Gordon, D.F.& Ridgway, E.C. (1993). A thyrotrope-specific variant of Pit-1 transactivates the thyrotropin beta promoter. *J Biol Chem* , 268 , 20818-20824.
180. Haugen, B.R., Gordon, D.F., Nelson, A.R., Wood, W.M.& Ridgway, E.C. (1994). The combination of Pit-1 and Pit-1T have a synergistic stimulatory effect on the thyrotropin beta-subunit promoter but not the growth hormone or prolactin promoters. *Mol Endocrinol* , 8 , 1574-1582.
181. Flynn, T.R., Hollenberg, A.N., Cohen, O., Menke, J.B., Usala, S.J., Tollin, S., Hegarty, M.K.& Wondisford, F.E. (1994). A novel C-terminal domain in the thyroid hormone receptor selectively mediates thyroid hormone inhibition. *J Biol Chem* , 269 , 32713-32716.
182. Haugen, B.R., McDermott, M.T., Gordon, D.F., Rupp, C.L., Wood, W.M.& Ridgway, E.C. (1996). Determinants of thyrotrope-specific thyrotropin beta promoter activation. Cooperation of Pit-1 with another factor. *J Biol Chem* , 271 , 385-389.
183. Gordon, D.F., Lewis, S.R., Haugen, B.R., James, R.A., McDermott, M.T., Wood, W.M.& Ridgway, E.C. (1997). Pit-1 and GATA-2 interact and functionally cooperate to activate the thyrotropin beta-subunit promoter. *J Biol Chem* , 272 , 24339-24347.
184. Bach, I., Rhodes, S.J., Pearse, R.V., 2nd, Heinzel, T., Gloss, B., Scully, K.M., Sawchenko, P.E.& Rosenfeld, M.G. (1995). P-Lim, a LIM homeodomain factor, is expressed during pituitary organ and cell commitment and synergizes with Pit-1. *Proc Natl Acad Sci U S A* , 92 , 2720-2724.
185. Deladoey, J., Fluck, C., Buyukgebiz, A., Kuhlmann, B.V., Eble, A., Hindmarsh, P.C., Wu, W.& Mullis, P.E. (1999). "Hot spot" in the PROP1 gene responsible for combined pituitary hormone deficiency. *J Clin Endocrinol Metab* , 84 , 1645-1650.
186. Vieira, T.C., Dias da Silva, M.R., Cerutti, J.M., Brunner, E., Borges, M., Arnaldi, L.T., Kopp, P.& Abucham, J. (2003). Familial combined pituitary hormone deficiency due to a novel mutation R99Q in the hot spot region of Prophet of Pit-1 presenting as constitutional growth delay. *J Clin Endocrinol Metab* , 88 , 38-44.
187. Re, R.N., Kourides, I.A., Ridgway, E.C., Weintraub, B.D.& Maloof, F. (1976). The effect of glucocorticoid administration on human pituitary secretion of thyrotropin and prolactin. *J Clin Endocrinol Metab* , 43 , 338-346.
188. Beck-Peccoz, P., Brucker-Davis, F., Persani, L., Smallridge, R.C.& Weintraub, B.D. (1996). Thyrotropin-secreting pituitary tumors. *Endocr Rev* , 17 , 610-638.
189. Ross, D.S., Ellis, M.F., Milbury, P.& Ridgway, E.C. (1987). A comparison of changes in plasma thyrotropin beta- and alpha-subunits, and mouse thyrotropic tumor thyrotropin beta- and alpha-subunit mRNA concentrations after in vivo dexamethasone or T3 administration. *Metabolism* , 36 , 799-803.
190. Coiro, V., Volpi, R., Cataldo, S., Capretti, L., Caffarri, G., Pilla, S.& Chiodera, P. (2000). Dopaminergic and cholinergic involvement in the inhibitory effect of dexamethasone on the TSH response to TRH. *J Investig Med* , 48 , 133-136.
191. Ahlquist, J.A., Franklyn, J.A., Wood, D.F., Balfour, N.J., Docherty, K., Sheppard, M.C.& Ramsden, D.B. (1987). Hormonal regulation of thyrotrophin synthesis and secretion. *Horm Metab Res Suppl* , 17 , 86-89.
192. Glass, C.K., Holloway, J.M., Devary, O.V.& Rosenfeld, M.G. (1988). The thyroid hormone receptor binds with opposite transcriptional effects to a common sequence motif in thyroid hormone

and estrogen response elements. *Cell* , 54 , 313-323.

193. Lamberts, S.W., Zuyderwijk, J., den Holder, F., van Koetsveld, P.& Hofland, L. (1989). Studies on the conditions determining the inhibitory effect of somatostatin on adrenocorticotropin, prolactin and thyrotropin release by cultured rat pituitary cells. *Neuroendocrinology* , 50 , 44-50.
194. Weeke, J., Hansen, A.P.& Lundaek, K. (1975). Inhibition by somatostatin of basal levels of serum thyrotropin (TSH) in normal men. *J Clin Endocrinol Metab* , 41 , 168-171.
195. Weeke, J., Christensen, S.E., Hansen, A.P., Laurberg, P.& Lundbaek, K. (1980). Somatostatin and the 24 h levels of serum TSH, T<sub>3</sub>, T<sub>4</sub>, and reverse T<sub>3</sub> in normals, diabetics and patients treated for myxoedema. *Acta Endocrinol (Copenh)* , 94 , 30-37.
196. Arimura, A.& Schally, A.V. (1976). Increase in basal and thyrotropin-releasing hormone (TRH)-stimulated secretion of thyrotropin (TSH) by passive immunization with antiserum to somatostatin in rats. *Endocrinology* , 98 , 1069-1072.
197. Yang, I., Woo, J., Kim, S., Kim, J., Kim, Y.& Choi, Y. (1996). Suppression of TRH-stimulated TSH secretion by glucose-induced hypothalamic somatostatin release. *Horm Metab Res* , 28 , 553-557.
198. Comi, R.J., Gesundheit, N., Murray, L., Gorden, P.& Weintraub, B.D. (1987). Response of thyrotropin-secreting pituitary adenomas to a long-acting somatostatin analogue. *N Engl J Med* , 317 , 12-17.
199. Beck-Peccoz, P., Mariotti, S., Guillausseau, P.J., Medri, G., Piscitelli, G., Bertoli, A., Barbarino, A., Rondena, M., Chanson, P., Pinchera, A.& et al. (1989). Treatment of hyperthyroidism due to inappropriate secretion of thyrotropin with the somatostatin analog SMS 201-995. *J Clin Endocrinol Metab* , 68 , 208-214.
200. Reisine, T.& Bell, G.I. (1995). Molecular biology of somatostatin receptors. *Endocr Rev* , 16 , 427-442.
201. Nabekura, J., Mizuno, Y.& Oomura, Y. (1989). Inhibitory effect of somatostatin on vagal motoneurons in the rat brain stem in vitro. *Am J Physiol* , 256 , C155-159.
202. Nilsson, T., Arkhammar, P., Rorsman, P.& Berggren, P.O. (1989). Suppression of insulin release by galanin and somatostatin is mediated by a G-protein. An effect involving repolarization and reduction in cytoplasmic free Ca<sup>2+</sup> concentration. *J Biol Chem* , 264 , 973-980.
203. De Groef, B., Goris, N., Arckens, L., Kuhn, E.R.& Darras, V.M. (2003). Corticotropin-Releasing Hormone (CRH)-Induced Thyrotropin Release is Directly Mediated Through CRH Receptor Type 2 on Thyrotropes. *Endocrinology* , 144 , 5537-5544.
204. De Groef, B., Geris, K.L., Manzano, J., Bernal, J., Millar, R.P., Abou-Samra, A.B., Porter, T.E., Iwasawa, A., Kuhn, E.R.& Darras, V.M. (2003). Involvement of thyrotropin-releasing hormone receptor, somatostatin receptor subtype 2 and corticotropin-releasing hormone receptor type 1 in the control of chicken thyrotropin secretion. *Mol Cell Endocrinol* , 203 , 33-39.
205. Kennedy, A.R., Todd, J.F., Stanley, S.A., Abbott, C.R., Small, C.J., Ghatei, M.A.& Bloom, S.R. (2000). Melanin-concentrating hormone (MCH) suppresses thyroid stimulating hormone (TSH) release, in vivo and in vitro, via the hypothalamus and the pituitary. *Endocrinology* , 142 , 3265-3268.
206. Fukuda, H.& Greer, M.A. (1975). The effect of basal hypothalamic deafferentation on the nycthemeral rhythm of plasma TSH. *Endocrinology* , 97 , 749-752.
207. Spaulding, S.W., Burrow, G.N., Donabedian, R.& van Woert, M. (1972). L-DOPA suppression of thyrotropin-releasing hormone response in man. *J Clin Endocrinol Metab* , 35 , 182-185.

208. Besses, G.S., Burrow, G.N., Spaulding, S.W.& Donabedian, R.K. (1975). Dopamine infusion acutely inhibits the TSH and prolactin response to TRH. *J Clin Endocrinol Metab* , 41 , 985-988.
209. Leebaw, W.F., Lee, L.A.& Woolf, P.D. (1978). Dopamine affects basal and augmented pituitary hormone secretion. *J Clin Endocrinol Metab* , 47 , 480-487.
210. Lewis, B.M., Dieguez, C., Lewis, M.D.& Scanlon, M.F. (1987). Dopamine stimulates release of thyrotrophin-releasing hormone from perfused intact rat hypothalamus via hypothalamic D2-receptors. *J Endocrinol* , 115 , 419-424.
211. Lewis, B.M., Dieguez, C., Ham, J., Page, M.D., Creagh, F.M., Peters, J.R.& Scanlon, M.F. (1989). Effects of Glucose on Thyrotrophin-Releasing Hormone, Growth Hormone-Releasing Hormone, Somatostatin and Luteinizing Hormone-Releasing Hormone Release from Rat Hypothalamus in vitro. *J Neuroendocrinol* , 1 , 437-441.
212. Krulich, L., Giachetti, A., Marchlewska-Koj, A., Hefco, E.& Jameson, H.E. (1977). On the role of the central noradrenergic and dopaminergic systems in the regulation of TSH secretion in the rat. *Endocrinology* , 100 , 496-505.
213. Krulich, L., Mayfield, M.A., Steele, M.K., McMillen, B.A., McCann, S.M.& Koenig, J.I. (1982). Differential effects of pharmacological manipulations of central alpha 1- and alpha 2-adrenergic receptors on the secretion of thyrotropin and growth hormone in male rats. *Endocrinology* , 110 , 796-804.
214. Gershengorn, M.C., Rebecchi, M.J., Geras, E.& Arevalo, C.O. (1980). Thyrotropin-releasing hormone (TRH) action in mouse thyrotropic tumor cells in culture: evidence against a role for adenosine 3',5'-monophosphate as a mediator of TRH-stimulated thyrotropin release. *Endocrinology* , 107 , 665-670.
215. Dieguez, C., Foord, S.M., Peters, J.R., Hall, R.& Scanlon, M.F. (1985). Alpha 1-adrenoreceptors and alpha 1-adrenoreceptor-mediated thyrotropin release in cultures of euthyroid and hypothyroid rat anterior pituitary cells. *Endocrinology* , 117 , 624-630.
216. Sharp, B., Morley, J.E., Carlson, H.E., Gordon, J., Briggs, J., Melmed, S.& Hershman, J.M. (1981). The role of opiates and endogenous opioid peptides in the regulation of rat TSH secretion. *Brain Res* , 219 , 335-344.
217. Samuels, M.H., Kramer, P., Wilson, D.& Sexton, G. (1994). Effects of naloxone infusions on pulsatile thyrotropin secretion. *J Clin Endocrinol Metab* , 78 , 1249-1252.
218. Peuranen, E., Vasar, E., Koks, S., Volke, V., Lang, A., Rauhala, P.& Mannisto, P.T. (1995). Further studies on the role of cholecystokinin-A and B receptors in secretion of anterior pituitary hormones in male rats. *Neuropeptides* , 28 , 1-11.
219. Santos, C.V., Pazos-Moura, C.C.& Moura, E.G. (1995). Effect of gastrin-releasing peptide (GRP) and GRP antagonists on TSH secretion from rat isolated pituitaries. *Life Sci* , 57 , 911-915.
220. Fekete, C., Kelly, J., Mihaly, E., Sarkar, S., Rand, W.M., Legradi, G., Emerson, C.H.& Lechan, R.M. (2001). Neuropeptide Y has a central inhibitory action on the hypothalamic- pituitary-thyroid axis. *Endocrinology* , 142 , 2606-2613.
221. Ciosek, J.& Stempniak, B. (1997). The influence of vasopressin or oxytocin on thyroid-stimulating hormone and thyroid hormones' concentrations in blood plasma of euthyroid rats. *J Physiol Pharmacol* , 48 , 813-823.
222. Beak, S.A., Small, C.J., Illovaiskaia, I., Hurley, J.D., Ghatei, M.A., Bloom, S.R.& Smith, D.M. (1996). Glucagon-like peptide-1 (GLP-1) releases thyrotropin (TSH): characterization of binding sites

- for GLP-1 on alpha-TSH cells. *Endocrinology* , 137 , 4130-4138.
223. Arvat, E., Gianotti, L., Ramunni, J., Grottoli, S., Brossa, P.C., Bertagna, A., Camanni, F.& Ghigo, E. (1995). Effect of galanin on basal and stimulated secretion of prolactin, gonadotropins, thyrotropin, adrenocorticotropin and cortisol in humans. *Eur J Endocrinol* , 133 , 300-304.
224. Nillni, E.A., Vaslet, C., Harris, M., Hollenberg, A., Bjorbak, C.& Flier, J.S. (2000). Leptin regulates prothyrotropin-releasing hormone biosynthesis. Evidence for direct and indirect pathways. *J Biol Chem* , 275 , 36124-36133.
225. Seoane, L.M., Carro, E., Tovar, S., Casanueva, F.F.& Dieguez, C. (2000). Regulation of in vivo TSH secretion by leptin. *Regul Pept* , 92 , 25-29.
226. Flier, J.S., Harris, M.& Hollenberg, A.N. (2000). Leptin, nutrition, and the thyroid: the why, the wherefore, and the wiring. *J Clin Invest* , 105 , 859-861.
227. Ozawa, M., Sato, K., Han, D.C., Kawakami, M., Tsushima, T.& Shizume, K. (1988). Effects of tumor necrosis factor-alpha/cachectin on thyroid hormone metabolism in mice. *Endocrinology* , 123 , 1461-1467.
228. Pang, X.P., Hershman, J.M., Mirell, C.J.& Pekary, A.E. (1989). Impairment of hypothalamic-pituitary-thyroid function in rats treated with human recombinant tumor necrosis factor-alpha (cachectin). *Endocrinology* , 125 , 76-84.
229. Dubuis, J.M., Dayer, J.M., Siegrist-Kaiser, C.A.& Burger, A.G. (1988). Human recombinant interleukin-1 beta decreases plasma thyroid hormone and thyroid stimulating hormone levels in rats. *Endocrinology* , 123 , 2175-2181.
230. van der Poll, T., Romijn, J.A., Wiersinga, W.M.& Sauerwein, H.P. (1990). Tumor necrosis factor: a putative mediator of the sick euthyroid syndrome in man. *J Clin Endocrinol Metab* , 71 , 1567-1572.
231. Wassen, F.W., Moerings, E.P., Van Toor, H., De Vrey, E.A., Hennemann, G.& Everts, M.E. (1996). Effects of interleukin-1 beta on thyrotropin secretion and thyroid hormone uptake in cultured rat anterior pituitary cells. *Endocrinology* , 137 , 1591-1598.
232. Melmed, S. (1998). gp130-related cytokines and their receptors in the pituitary. *TEM* , 9 , 155.
233. Rettori, V., Dees, W.L., Hiney, J.K., Lyson, K.& McCann, S.M. (1994). An interleukin-1-alpha-like neuronal system in the preoptic-hypothalamic region and its induction by bacterial lipopolysaccharide in concentrations which alter pituitary hormone release. *Neuroimmunomodulation* , 1 , 251-258.
234. Koenig, J.I., Snow, K., Clark, B.D., Toni, R., Cannon, J.G., Shaw, A.R., Dinarello, C.A., Reichlin, S., Lee, S.L.& Lechan, R.M. (1990). Intrinsic pituitary interleukin-1 beta is induced by bacterial lipopolysaccharide. *Endocrinology* , 126 , 3053-3058.
235. van Haasteren, G.A., van der Meer, M.J., Hermus, A.R., Linkels, E., Klootwijk, W., Kaptein, E., van Toor, H., Sweep, C.G., Visser, T.J.& de Greef, W.J. (1994). Different effects of continuous infusion of interleukin-1 and interleukin-6 on the hypothalamic-hypophysial-thyroid axis. *Endocrinology* , 135 , 1336-1345.
236. Bartalena, L., Grasso, L., Brogioni, S.& Martino, E. (1994). Interleukin 6 effects on the pituitary-thyroid axis in the rat. *Eur J Endocrinol* , 131 , 302-306.
237. Kakucska, I., Romero, L.I., Clark, B.D., Rondeel, J.M., Qi, Y., Alex, S., Emerson, C.H.& Lechan, R.M. (1994). Suppression of thyrotropin-releasing hormone gene expression by interleukin-1-beta in the rat: implications for nonthyroidal illness. *Neuroendocrinology* , 59 , 129-137.

238. Baur, A., Bauer, K., Jarry, H.& Kohrle, J. (2000). Effects of proinflammatory cytokines on anterior pituitary 5'- deiodinase type I and type II. *J Endocrinol* , 167 , 505-515.
239. Spangelo, B.L., MacLeod, R.M.& Isakson, P.C. (1990). Production of interleukin-6 by anterior pituitary cells in vitro. *Endocrinology* , 126 , 582-586.
240. Vankelecom, H., Carmeliet, P., Van Damme, J., Billiau, A.& Denef, C. (1989). Production of interleukin-6 by folliculo-stellate cells of the anterior pituitary gland in a histiotypic cell aggregate culture system. *Neuroendocrinology* , 49 , 102-106.
241. Akieda-Asai, S., Zaima, N., Ikegami, K., Kahyo, T., Yao, I., Hatanaka, T., Iemura, S., Sugiyama, R., Yokozeki, T., Eishi, Y., Koike, M., Ikeda, K., Chiba, T., Yamaza, H., Shimokawa, I., Song, S.Y., Matsuno, A., Mizutani, A., Sawabe, M., Chao, M.V., Tanaka, M., Kanaho, Y., Natsume, T., Sugimura, H., Date, Y., McBurney, M.W., Guarente, L.& Setou, M. (2010). SIRT1 Regulates Thyroid-Stimulating Hormone Release by Enhancing PIP5Kgamma Activity through Deacetylation of Specific Lysine Residues in Mammals. *PLoS One* , 5 , e11755.
242. Prummel, M.F., Brokken, L.J.& Wiersinga, W.M. (2004). Ultra short-loop feedback control of thyrotropin secretion. *Thyroid* , 14 , 825-829.
243. Bockmann, J., Winter, C., Wittkowski, W., Kreutz, M.R.& Bockers, T.M. (1997). Cloning and expression of a brain-derived TSH receptor. *Biochem Biophys Res Commun* , 238 , 173-178.
244. Crisanti, P., Omri, B., Hughes, E., Meduri, G., Hery, C., Clauser, E., Jacquemin, C.& Saunier, B. (2001). The expression of thyrotropin receptor in the brain. *Endocrinology* , 142 , 812-822.
245. Prummel, M.F., Brokken, L.J., Meduri, G., Misrahi, M., Bakker, O.& Wiersinga, W.M. (2000). Expression of the thyroid-stimulating hormone receptor in the folliculo- stellate cells of the human anterior pituitary. *J Clin Endocrinol Metab* , 85 , 4347-4353.
246. Wartofsky, L.& Dickey, R.A. (2005). The evidence for a narrower thyrotropin reference range is compelling. *J Clin Endocrinol Metab* , 90 , 5483-5488.
247. Surks, M.I., Goswami, G.& Daniels, G.H. (2005). The thyrotropin reference range should remain unchanged. *J Clin Endocrinol Metab* , 90 , 5489-5496.
248. Surks, M.I.& Boucail, L. (2010). Age- and race-based serum thyrotropin reference limits. *J Clin Endocrinol Metab* , 95 , 496-502.
249. Surks, M.I.& Hollowell, J.G. (2007). Age-specific distribution of serum thyrotropin and antithyroid antibodies in the US population: implications for the prevalence of subclinical hypothyroidism. *J Clin Endocrinol Metab* , 92 , 4575-4582.
250. Mariotti, S., Franceschi, C., Cossarizza, A.& Pinchera, A. (1995). The aging thyroid. *Endocr Rev* , 16 , 686-715.
251. Hansen, P.S., Brix, T.H., Sorensen, T.I., Kyvik, K.O.& Hegedus, L. (2004). Major genetic influence on the regulation of the pituitary-thyroid axis: a study of healthy Danish twins. *J Clin Endocrinol Metab* , 89 , 1181-1187.
252. Mariotti, S., Naitza, S.& Cao, A. (2010). Phosphodiesterase 8B (PDE8B) gene variants and TSH levels. *Hot Thyroidtology* , 3 , [http://www.hotthyroidology.com/editorial\\_211.html](http://www.hotthyroidology.com/editorial_211.html).
253. Arnaud-Lopez, L., Usala, G., Ceresini, G., Mitchell, B.D., Pilia, M.G., Piras, M.G., Sestu, N., Maschio, A., Busonero, F., Albai, G., Dei, M., Lai, S., Mulas, A., Crisponi, L., Tanaka, T., Bandinelli, S., Guralnik, J.M., Loi, A., Balaci, L., Sole, G., Prinzis, A., Mariotti, S., Shuldiner, A.R., Cao, A., Schlessinger, D., Uda, M., Abecasis, G.R., Nagaraja, R., Sanna, S.& Naitza, S. (2008).

Phosphodiesterase 8B gene variants are associated with serum TSH levels and thyroid function. Am J Hum Genet , 82 , 1270-1280.

254. Kourides, I.A., Ridgway, E.C., Weintraub, B.D., Bigos, S.T., Gershengorn, M.C.& Maloof, F. (1977). Thyrotropin-induced hyperthyroidism: use of alpha and beta subunit levels to identify patients with pituitary tumors. J Clin Endocrinol Metab , 45 , 534-543.
255. Kuzuya, N., Inoue, K., Ishibashi, M., Murayama, Y., Koide, Y., Ito, K., Yamaji, T.& Yamashita, K. (1990). Endocrine and immunohistochemical studies on thyrotropin (TSH)-secreting pituitary adenomas: responses of TSH, alpha-subunit, and growth hormone to hypothalamic releasing hormones and their distribution in adenoma cells. J Clin Endocrinol Metab , 71 , 1103-1111.
256. Gesundheit, N., Petrick, P.A., Nissim, M., Dahlberg, P.A., Doppman, J.L., Emerson, C.H., Braverman, L.E., Oldfield, E.H.& Weintraub, B.D. (1989). Thyrotropin-secreting pituitary adenomas: clinical and biochemical heterogeneity. Case reports and follow-up of nine patients. Ann Intern Med , 111 , 827-835.
257. Faglia, G., Beck-Peccoz, P., Piscitelli, G.& Medri, G. (1987). Inappropriate secretion of thyrotropin by the pituitary. Horm Res , 26 , 79-99.
258. Spencer, C.A., Lai-Rosenfeld, A.O., Guttler, R.B., LoPresti, J., Marcus, A.O., Nimalasuriya, A., Eigen, A., Doss, R.C., Green, B.J.& Nicoloff, J.T. (1986). Thyrotropin secretion in thyrotoxic and thyroxine-treated patients: assessment by a sensitive immunoenzymometric assay. J Clin Endocrinol Metab , 63 , 349-355.
259. Spencer, C.A., LoPresti, J.S., Patel, A., Guttler, R.B., Eigen, A., Shen, D., Gray, D.& Nicoloff, J.T. (1990). Applications of a new chemiluminometric thyrotropin assay to subnormal measurement. J Clin Endocrinol Metab , 70 , 453-460.
260. Spencer, C.A., Schwarzbein, D., Guttler, R.B., LoPresti, J.S.& Nicoloff, J.T. (1993). Thyrotropin (TSH)-releasing hormone stimulation test responses employing third and fourth generation TSH assays. J Clin Endocrinol Metab , 76 , 494-498.
261. Greenspan, S.L., Klibanski, A., Schoenfeld, D.& Ridgway, E.C. (1986). Pulsatile secretion of thyrotropin in man. J Clin Endocrinol Metab , 63 , 661-668.
262. Brabant, G., Brabant, A., Ranft, U., Ocran, K., Kohrle, J., Hesch, R.D.& von zur Muhlen, A. (1987). Circadian and pulsatile thyrotropin secretion in euthyroid man under the influence of thyroid hormone and glucocorticoid administration. J Clin Endocrinol Metab , 65 , 83-88.
263. Brabant, G., Prank, K., Ranft, U., Schuermeyer, T., Wagner, T.O., Hauser, H., Kummer, B., Feistner, H., Hesch, R.D.& von zur Muhlen, A. (1990). Physiological regulation of circadian and pulsatile thyrotropin secretion in normal man and woman. J Clin Endocrinol Metab , 70 , 403-409.
264. Samuels, M.H., Veldhuis, J.& Ridgway, E.C. (1995). Copulsatile release of thyrotropin and prolactin in normal and hypothyroid subjects. Thyroid , 5 , 369-372.
265. Weeke, J. (1973). Circadian variation of the serum thyrotropin level in normal subjects. Scand J Clin Lab Invest , 31 , 337-342.
266. Bartalena, L., Martino, E., Brandi, L.S., Falcone, M., Pacchiarotti, A., Ricci, C., Bogazzi, F., Grasso, L., Mammoli, C.& Pinchera, A. (1990). Lack of nocturnal serum thyrotropin surge after surgery. J Clin Endocrinol Metab , 70 , 293-296.
267. Parker, D.C., Pekary, A.E.& Hershman, J.M. (1976). Effect of normal and reversed sleep-wake cycles upon nyctohemeral rhythmicity of plasma thyrotropin: evidence suggestive of an inhibitory influence in sleep. J Clin Endocrinol Metab , 43 , 318-329.

268. Parker, D.C., Rossman, L.G., Pekary, A.E.& Hershman, J.M. (1987). Effect of 64-hour sleep deprivation on the circadian waveform of thyrotropin (TSH): further evidence of sleep-related inhibition of TSH release. *J Clin Endocrinol Metab* , 64 , 157-161.
269. Behrends, J., Prank, K., Dogu, E.& Brabant, G. (1998). Central nervous system control of thyrotropin secretion during sleep and wakefulness. *Horm Res* , 49 , 173-177.
270. Hirshberg, B., Veldhuis, J.D.& Sarlis, N.J. (2000). Diurnal thyrotropin secretion in short-term profound primary hypothyroidism: does it ever persist? *Thyroid* , 10 , 1101-1106.
271. Olsen, T., Laurberg, P.& Weeke, J. (1978). Low serum triiodothyronine and high serum reverse triiodothyronine in old age: an effect of disease not age. *J Clin Endocrinol Metab* , 47 , 1111-1115.
272. Sawin, C.T., Geller, A., Kaplan, M.M., Bacharach, P., Wilson, P.W.& Hershman, J.M. (1991). Low serum thyrotropin (thyroid-stimulating hormone) in older persons without hyperthyroidism. *Arch Intern Med* , 151 , 165-168.
273. Sawin, C.T., Geller, A., Wolf, P.A., Belanger, A.J., Baker, E., Bacharach, P., Wilson, P.W., Benjamin, E.J.& D'Agostino, R.B. (1994). Low serum thyrotropin concentrations as a risk factor for atrial fibrillation in older persons. *N Engl J Med* , 331 , 1249-1252.
274. Mariotti, S., Barbesino, G., Caturegli, P., Bartalena, L., Sansoni, P., Fagnoni, F., Monti, D., Fagiolo, U., Franceschi, C.& Pinchera, A. (1993). Complex alteration of thyroid function in healthy centenarians. *J Clin Endocrinol Metab* , 77 , 1130-1134.
275. Cuttelod, S., Lemarchand-Beraud, T., Magnenat, P., Perret, C., Poli, S.& Vannotti, A. (1974). Effect of age and role of kidneys and liver on thyrotropin turnover in man. *Metabolism* , 23 , 101-113.
276. van Coevorden, A., Laurent, E., Decoster, C., Kerkhofs, M., Neve, P., van Cauter, E.& Mockel, J. (1989). Decreased basal and stimulated thyrotropin secretion in healthy elderly men. *J Clin Endocrinol Metab* , 69 , 177-185.
277. Balsam, A.& Ingbar, S.H. (1979). Observations on the factors that control the generation of triiodothyronine from thyroxine in rat liver and the nature of the defect induced by fasting. *J Clin Invest* , 63 , 1145-1156.
278. Harris, A.R., Fang, S.L., Hinerfeld, L., Braverman, L.E.& Vagenakis, A.G. (1979). The role of sulphydryl groups on the impaired hepatic 3',3,5-triiodothyronine generation from thyroxine in the hypothyroid, starved, fetal, and neonatal rodent. *J Clin Invest* , 63 , 516-524.
279. Kaplan, M.M. (1979). Subcellular alterations causing reduced hepatic thyroxine-5'-monodeiodinase activity in fasted fats. *Endocrinology* , 104 , 58-64.
280. Kaplan, M.M., Tatro, J.B., Breitbart, R.& Larsen, P.R. (1979). Comparison of thyroxine and 3,3',5'-triiodothyronine metabolism in rat kidney and liver homogenates. *Metabolism* , 28 , 1139-1146.
281. Kinlaw, W.B., Schwartz, H.L.& Oppenheimer, J.H. (1985). Decreased serum triiodothyronine in starving rats is due primarily to diminished thyroidal secretion of thyroxine. *J Clin Invest* , 75 , 1238-1241.
282. Wartofsky, L.& Burman, K.D. (1982). Alterations in thyroid function in patients with systemic illness: the "euthyroid sick syndrome". *Endocr Rev* , 3 , 164-217.
283. Wehmann, R.E., Gregerman, R.I., Burns, W.H., Saral, R.& Santos, G.W. (1985). Suppression of thyrotropin in the low-thyroxine state of severe nonthyroidal illness. *N Engl J Med* , 312 , 546-552.
284. Hugues, J.N., Burger, A.G., Pekary, A.E.& Hershman, J.M. (1984). Rapid adaptations of serum thyrotrophin, triiodothyronine and reverse triiodothyronine levels to short-term starvation and

- refeeding. *Acta Endocrinol (Copenh)* , 105 , 194-199.
285. Samuels, M.H.& Kramer, P. (1996). Differential effects of short-term fasting on pulsatile thyrotropin, gonadotropin, and alpha-subunit secretion in healthy men—a clinical research center study. *J Clin Endocrinol Metab* , 81 , 32-36.
286. Samuels, M.H.& Kramer, P. (1996). Effects of metoclopramide on fasting-induced TSH suppression. *Thyroid* , 6 , 85-89.
287. Carlson, H.E., Drenick, E.J., Chopra, I.J.& Hershman, J.M. (1977). Alterations in basal and TRH-stimulated serum levels of thyrotropin, prolactin, and thyroid hormones in starved obese men. *J Clin Endocrinol Metab* , 45 , 707-713.
288. Spencer, C.A., Lum, S.M., Wilber, J.F., Kaptein, E.M.& Nicoloff, J.T. (1983). Dynamics of serum thyrotropin and thyroid hormone changes in fasting. *J Clin Endocrinol Metab* , 56 , 883-888.
289. Shi, Z.X., Levy, A.& Lightman, S.L. (1993). The effect of dietary protein on thyrotropin-releasing hormone and thyrotropin gene expression. *Brain Res* , 606 , 1-4.
290. Blake, N.G., Eckland, D.J., Foster, O.J.& Lightman, S.L. (1991). Inhibition of hypothalamic thyrotropin-releasing hormone messenger ribonucleic acid during food deprivation. *Endocrinology* , 129 , 2714-2718.
291. Hugues, J.N., Enjalbert, A., Moyse, E., Shu, C., Voirol, M.J., Sebaoun, J.& Epelbaum, J. (1986). Differential effects of passive immunization with somatostatin antiserum on adenohypophysial hormone secretions in starved rats. *J Endocrinol* , 109 , 169-174.
292. Kok, P., Roelfsema, F., Langendonk, J.G., Frolich, M., Burggraaf, J., Meinders, A.E.& Pijl, H. (2005). High circulating thyrotropin levels in obese women are reduced after body weight loss induced by caloric restriction. *J Clin Endocrinol Metab* , 90 , 4659-4663.
293. Kok, P., Roelfsema, F., Frolich, M., Meinders, A.E.& Pijl, H. (2005). Spontaneous Diurnal Tsh Secretion Is Enhanced in Proportion to Circulating Leptin in Obese Premenopausal Women. *J Clin Endocrinol Metab* ,
294. Legradi, G., Emerson, C.H., Ahima, R.S., Flier, J.S.& Lechan, R.M. (1997). Leptin prevents fasting-induced suppression of prothyrotropin-releasing hormone messenger ribonucleic acid in neurons of the hypothalamic paraventricular nucleus. *Endocrinology* , 138 , 2569-2576.
295. Harris, M., Aschkenasi, C., Elias, C.F., Chandrankunnel, A., Nilni, E.A., Bjorbaek, C., Elmquist, J.K., Flier, J.S.& Hollenberg, A.N. (2001). Transcriptional regulation of the thyrotropin-releasing hormone gene by leptin and melanocortin signaling. *J Clin Invest* , 107 , 111-120.
296. Kim, M.S., Small, C.J., Stanley, S.A., Morgan, D.G., Seal, L.J., Kong, W.M., Edwards, C.M., Abusnana, S., Sunter, D., Ghatei, M.A.& Bloom, S.R. (2000). The central melanocortin system affects the hypothalamo-pituitary thyroid axis and may mediate the effect of leptin. *J Clin Invest* , 105 , 1005-1011.
297. Guo, F., Bakal, K., Minokoshi, Y.& Hollenberg, A.N. (2004). Leptin signaling targets the thyrotropin-releasing hormone gene promoter in vivo. *Endocrinology* , 145 , 2221-2227.
298. Huo, L., Munzberg, H., Nilni, E.A.& Bjorbaek, C. (2004). Role of signal transducer and activator of transcription 3 in regulation of hypothalamic trh gene expression by leptin. *Endocrinology* , 145 , 2516-2523.
299. Fekete, C., Legradi, G., Mihaly, E., Huang, Q.H., Tatro, J.B., Rand, W.M., Emerson, C.H.& Lechan, R.M. (2000). alpha-Melanocyte-stimulating hormone is contained in nerve terminals

- innervating thyrotropin-releasing hormone-synthesizing neurons in the hypothalamic paraventricular nucleus and prevents fasting-induced suppression of prothyrotropin-releasing hormone gene expression. *J Neurosci* , 20 , 1550-1558.
300. Mihaly, E., Fekete, C., Tatro, J.B., Liposits, Z., Stopa, E.G.& Lechan, R.M. (2000). Hypophysiotropic thyrotropin-releasing hormone-synthesizing neurons in the human hypothalamus are innervated by neuropeptide Y, agouti-related protein, and alpha-melanocyte-stimulating hormone. *J Clin Endocrinol Metab* , 85 , 2596-2603.
301. Fekete, C., Marks, D.L., Sarkar, S., Emerson, C.H., Rand, W.M., Cone, R.D.& Lechan, R.M. (2004). Effect of Agouti-related protein in regulation of the hypothalamic-pituitary-thyroid axis in the melanocortin 4 receptor knockout mouse. *Endocrinology* , 145 , 4816-4821.
302. Fekete, C., Sarkar, S., Rand, W.M., Harney, J.W., Emerson, C.H., Bianco, A.C., Beck-Sickinger, A.& Lechan, R.M. (2002). Neuropeptide Y1 and Y5 receptors mediate the effects of neuropeptide Y on the hypothalamic-pituitary-thyroid axis. *Endocrinology* , 143 , 4513-4519.
303. Wittmann, G., Liposits, Z., Lechan, R.M.& Fekete, C. (2002). Medullary adrenergic neurons contribute to the neuropeptide Y-ergic innervation of hypophysiotropic thyrotropin-releasing hormone-synthesizing neurons in the rat. *Neurosci Lett* , 324 , 69-73.
304. Kaplan, M.M., Larsen, P.R., Crantz, F.R., Dzau, V.J., Rossing, T.H.& Haddow, J.E. (1982). Prevalence of abnormal thyroid function test results in patients with acute medical illnesses. *Am J Med* , 72 , 9-16.
305. Romijn, J.A., Adriaanse, R., Brabant, G., Prank, K., Endert, E.& Wiersinga, W.M. (1990). Pulsatile secretion of thyrotropin during fasting: a decrease of thyrotropin pulse amplitude. *J Clin Endocrinol Metab* , 70 , 1631-1636.
306. Kamat, V., Hecht, W.L.& Rubin, R.T. (1995). Influence of meal composition on the postprandial response of the pituitary-thyroid axis. *Eur J Endocrinol* , 133 , 75-79.
307. Oppert, J.M., Dussault, J.H., Tremblay, A., Despres, J.P., Theriault, G.& Bouchard, C. (1994). Thyroid hormones and thyrotropin variations during long term overfeeding in identical twins. *J Clin Endocrinol Metab* , 79 , 547-553.
308. Knudsen, N., Laurberg, P., Rasmussen, L.B., Bulow, I., Perrild, H., Ovesen, L.& Jorgensen, T. (2005). Small differences in thyroid function may be important for body mass index and the occurrence of obesity in the population. *J Clin Endocrinol Metab* , 90 , 4019-4024.
309. Ehrmann, D.A., Weinberg, M.& Sarne, D.H. (1989). Limitations to the use of a sensitive assay for serum thyrotropin in the assessment of thyroid status. *Arch Intern Med* , 149 , 369-372.
310. Faber, J., Kirkegaard, C., Rasmussen, B., Westh, H., Busch-Sorensen, M.& Jensen, I.W. (1987). Pituitary-thyroid axis in critical illness. *J Clin Endocrinol Metab* , 65 , 315-320.
311. Hamblin, P.S., Dyer, S.A., Mohr, V.S., Le Grand, B.A., Lim, C.F., Tuxen, D.V., Topliss, D.J.& Stockigt, J.R. (1986). Relationship between thyrotropin and thyroxine changes during recovery from severe hypothyroxinemia of critical illness. *J Clin Endocrinol Metab* , 62 , 717-722.
312. Romijn, J.A.& Wiersinga, W.M. (1990). Decreased nocturnal surge of thyrotropin in nonthyroidal illness. *J Clin Endocrinol Metab* , 70 , 35-42.
313. Custo, N., Scafidi, V., Gallo, S.& Notarbartolo, A. (1994). Deficient pulsatile thyrotropin secretion in the low-thyroid-hormone state of severe non-thyroidal illness. *Eur J Endocrinol* , 130 , 132-136.

314. Van den Berghe, G., de Zegher, F., Veldhuis, J.D., Wouters, P., Gouwy, S., Stockman, W., Weekers, F., Schetz, M., Lauwers, P., Bouillon, R.& Bowers, C.Y. (1997). Thyrotrophin and prolactin release in prolonged critical illness: dynamics of spontaneous secretion and effects of growth hormone-secretagogues. *Clin Endocrinol (Oxf)* , 47 , 599-612.
315. Bacci, V., Schussler, G.C.& Kaplan, T.B. (1982). The relationship between serum triiodothyronine and thyrotropin during systemic illness. *J Clin Endocrinol Metab* , 54 , 1229-1235.
316. Brent, G.A., Hershman, J.M.& Braunstein, G.D. (1986). Patients with severe nonthyroidal illness and serum thyrotropin concentrations in the hypothyroid range. *Am J Med* , 81 , 463-466.
317. Magner, J., Roy, P., Fainter, L., Barnard, V.& Fletcher, P., Jr. (1997). Transiently decreased sialylation of thyrotropin (TSH) in a patient with the euthyroid sick syndrome. *Thyroid* , 7 , 55-61.
318. De Groot, L.J. (1999). Dangerous dogmas in medicine: the nonthyroidal illness syndrome. *J Clin Endocrinol Metab* , 84 , 151-164.
319. Fliers, E., Alkemade, A.& Wiersinga, W.M. (2001). The hypothalamic-pituitary-thyroid axis in critical illness. *Best Pract Res Clin Endocrinol Metab* , 15 , 453-464.
320. Hennessey, J.& Jackson, I.M.D. (1996). The interface between thyroid hormones and psychiatry. *The endocrinologist* , 6 , 214.
321. Bartalena, L., Placidi, G.F., Martino, E., Falcone, M., Pellegrini, L., Dell'Osso, L., Pacchiarotti, A.& Pinchera, A. (1990). Nocturnal serum thyrotropin (TSH) surge and the TSH response to TSH-releasing hormone: dissociated behavior in untreated depressives. *J Clin Endocrinol Metab* , 71 , 650-655.
322. Gold, M.S., Pottash, A.L., Extein, I., Martin, D.M., Howard, E., Mueller, E.A., 3rd& Sweeney, D.R. (1981). The TRH test in the diagnosis of major and minor depression. *Psychoneuroendocrinology* , 6 , 159-169.
323. Chopra, I.J., Solomon, D.H.& Huang, T.S. (1990). Serum thyrotropin in hospitalized psychiatric patients: evidence for hyperthyrotropinemia as measured by an ultrasensitive thyrotropin assay. *Metabolism* , 39 , 538-543.
324. Loosen, P.T. (1988). Thyroid function in affective disorders and alcoholism. *Endocrinol Metab Clin North Am* , 17 , 55-82.
325. Banki, C.M., Bissette, G., Arato, M.& Nemeroff, C.B. (1988). Elevation of immunoreactive CSF TRH in depressed patients. *Am J Psychiatry* , 145 , 1526-1531.
326. Kirkegaard, C.& Carroll, B.J. (1980). Dissociation of TSH adrenocortical disturbances in endogenous depression. *Psychiatry Res* , 3 , 253-264.
327. O'Shanick, G.J.& Ellinwood, E.H., Jr. (1982). Persistent elevation of thyroid-stimulating hormone in women with bipolar affective disorder. *Am J Psychiatry* , 139 , 513-514.
328. Fliers, E., Guldhaar, S.E., Wiersinga, W.M.& Swaab, D.F. (1997). Decreased hypothalamic thyrotropin-releasing hormone gene expression in patients with nonthyroidal illness. *J Clin Endocrinol Metab* , 82 , 4032-4036.
329. Delitala, G., Tomasi, P.& Virdis, R. (1987). Prolactin, growth hormone and thyrotropin-thyroid hormone secretion during stress states in man. *Baillieres Clin Endocrinol Metab* , 1 , 391-414.
330. Chopra, I.J., Sakane, S.& Teco, G.N. (1991). A study of the serum concentration of tumor necrosis factor-alpha in thyroidal and nonthyroidal illnesses. *J Clin Endocrinol Metab* , 72 , 1113-1116.

331. Boelen, A., Platvoet-Ter Schiphorst, M.C.& Wiersinga, W.M. (1993). Association between serum interleukin-6 and serum 3,5,3'-triiodothyronine in nonthyroidal illness. *J Clin Endocrinol Metab* , 77 , 1695-1699.
332. Murai, H., Murakami, S., Ishida, K.& Sugawara, M. (1996). Elevated serum interleukin-6 and decreased thyroid hormone levels in postoperative patients and effects of IL-6 on thyroid cell function in vitro. *Thyroid* , 6 , 601-606.
333. Davies, P.H., Black, E.G., Sheppard, M.C.& Franklyn, J.A. (1996). Relation between serum interleukin-6 and thyroid hormone concentrations in 270 hospital in-patients with non-thyroidal illness. *Clin Endocrinol (Oxf)* , 44 , 199-205.
334. Hermus, R.M., Sweep, C.G., van der Meer, M.J., Ross, H.A., Smals, A.G., Benraad, T.J.& Kloppenborg, P.W. (1992). Continuous infusion of interleukin-1 beta induces a nonthyroidal illness syndrome in the rat. *Endocrinology* , 131 , 2139-2146.
335. Sweep, C.G., van der Meer, M.J., Ross, H.A., Vranckx, R., Visser, T.J.& Hermus, A.R. (1992). Chronic infusion of TNF-alpha reduces plasma T4 binding without affecting pituitary-thyroid activity in rats. *Am J Physiol* , 263 , E1099-1105.
336. Bartalena, L., Brogioni, S., Grasso, L.& Martino, E. (1995). Interleukin-6 and the thyroid. *Eur J Endocrinol* , 132 , 386-393.
337. Kaptein, E.M., Spencer, C.A., Kamiel, M.B.& Nicoloff, J.T. (1980). Prolonged dopamine administration and thyroid hormone economy in normal and critically ill subjects. *J Clin Endocrinol Metab* , 51 , 387-393.
338. Duick, D.S.& Wahner, H.W. (1979). Thyroid axis in patients with Cushing's syndrome. *Arch Intern Med* , 139 , 767-772.
339. Nicoloff, J.T., Fisher, D.A.& Appleman, M.D., Jr. (1970). The role of glucocorticoids in the regulation of thyroid function in man. *J Clin Invest* , 49 , 1922-1929.
340. Alkemade, A., Unmehopa, U.A., Wiersinga, W.M., Swaab, D.F.& Fliers, E. (2005). Glucocorticoids decrease thyrotropin-releasing hormone messenger ribonucleic acid expression in the paraventricular nucleus of the human hypothalamus. *J Clin Endocrinol Metab* , 90 , 323-327.
341. Gharib, H., Hodgson, S.F., Gastineau, C.F., Scholz, D.A.& Smith, L.A. (1972). Reversible hypothyroidism in Addison's disease. *Lancet* , 2 , 734-736.
342. Topliss, D.J., White, E.L.& Stockigt, J.R. (1980). Significance of thyrotropin excess in untreated primary adrenal insufficiency. *J Clin Endocrinol Metab* , 50 , 52-56.
343. Surks, M.I.& Sievert, R. (1995). Drugs and thyroid function. *N Engl J Med* , 333 , 1688-1694.
344. Faglia, G., Beck-Peccoz, P., Ferrari, C., Ambrosi, B., Spada, A.& Travaglini, P. (1973). Enhanced plasma thyrotrophin response to thyrotrophin-releasing hormone following oestradiol administration in man. *Clin Endocrinol (Oxf)* , 2 , 207-210.
345. Gershengorn, M.C., Marcus-Samuels, B.E.& Geras, E. (1979). Estrogens increase the number of thyrotropin-releasing hormone receptors on mammatropic cells in culture. *Endocrinology* , 105 , 171-176.
346. Morley, J.E., Sawin, C.T., Carlson, H.E., Longcope, C.& Hershman, J.M. (1981). The relationship of androgen to the thyrotropin and prolactin responses to thyrotropin-releasing hormone in hypogonadal and normal men. *J Clin Endocrinol Metab* , 52 , 173-176.
347. Arafah, B.M. (1994). Decreased levothyroxine requirement in women with hypothyroidism during

- androgen therapy for breast cancer. *Ann Intern Med* , 121 , 247-251.
348. Lippe, B.M., Van Herle, A.J., LaFranchi, S.H., Uller, R.P., Lavin, N.& Kaplan, S.A. (1975). Reversible hypothyroidism in growth hormone-deficient children treated with human growth hormone. *J Clin Endocrinol Metab* , 40 , 612-618.
349. Porter, B.A., Refetoff, S., Rosenfeld, R.L., De Groat, L.J., Lang, U.S.& Stark, O. (1973). Abnormal thyroxine metabolism in hyposomatrophic dwarfism and inhibition of responsiveness to TRH during GH therapy. *Pediatrics* , 51 , 668-674.
350. Jorgensen, J.O., Pedersen, S.A., Laurberg, P., Weeke, J., Skakkebaek, N.E.& Christiansen, J.S. (1989). Effects of growth hormone therapy on thyroid function of growth hormone-deficient adults with and without concomitant thyroxine-substituted central hypothyroidism. *J Clin Endocrinol Metab* , 69 , 1127-1132.
351. Trainer, P.J., Holly, J., Medbak, S., Rees, L.H.& Besser, G.M. (1994). The effect of recombinant IGF-I on anterior pituitary function in healthy volunteers. *Clin Endocrinol (Oxf)* , 41 , 801-807.
352. Little, K.Y., Garbutt, J.C., Mayo, J.P.& Mason, G. (1988). Lack of acute d-amphetamine effects on thyrotropin release. *Neuroendocrinology* , 48 , 304-307.
353. Rogol, A.D., Reeves, G.D., Varma, M.M.& Blizzard, R.M. (1979). Thyroid-stimulating hormone and prolactin responses to thyrotropin-releasing hormone during infusion of epinephrine and propranolol in man. *Neuroendocrinology* , 29 , 413-417.
354. Zgliczynski, S.& Kaniewski, M. (1980). Evidence for alpha-adrenergic receptors mediated TSH release in men. *Acta Endocrinol (Copenh)* , 95 , 172-176.
355. Valcavi, R., Dieguez, C., Azzarito, C., Artioli, C., Portioli, I.& Scanlon, M.F. (1987). Alpha-adrenoreceptor blockade with thymoxamine reduces basal thyrotrophin levels but does not influence circadian thyrotrophin changes in man. *J Endocrinol* , 115 , 187-191.
356. Lotti, G., Delitala, G., Devilla, L., Alagna, S.& Masala, A. (1977). Reduction of plasma triiodothyronine (T3) induced by propranolol. *Clin Endocrinol (Oxf)* , 6 , 405-410.
357. Plosker, S.M., Rabinovici, J., Montalvo, M.& Jaffe, R.B. (1995). Endogenous catecholamines suppress thyrotropin secretion during the early follicular phase of the menstrual cycle. *J Clin Endocrinol Metab* , 80 , 2530-2533.
358. Hay, I.D., Bayer, M.F., Kaplan, M.M., Klee, G.G., Larsen, P.R.& Spencer, C.A. (1991). American Thyroid Association assessment of current free thyroid hormone and thyrotropin measurements and guidelines for future clinical assays. The Committee on Nomenclature of the American Thyroid Association. *Clin Chem* , 37 , 2002-2008.
359. Baloch, Z., Carayon, P., Conte-Devolx, B., Demers, L.M., Feldt-Rasmussen, U., Henry, J.F., LiVosli, V.A., Niccoli-Sire, P., John, R., Ruf, J., Smyth, P.P., Spencer, C.A.& Stockigt, J.R. (2003). Laboratory medicine practice guidelines. Laboratory support for the diagnosis and monitoring of thyroid disease. *Thyroid* , 13 , 3-126.
360. Toft, A.D. (1988). Use of sensitive immunoradiometric assay for thyrotropin in clinical practice. *Mayo Clin Proc* , 63 , 1035-1042.
361. Martino, E., Bambini, G., Bartalena, L., Mammoli, C., Aghini-Lombardi, F., Baschieri, L.& Pinchera, A. (1986). Human serum thyrotrophin measurement by ultrasensitive immunoradiometric assay as a first-line test in the evaluation of thyroid function. *Clin Endocrinol (Oxf)* , 24 , 141-148.
362. Snyder, P.J.& Utiger, R.D. (1972). Response to thyrotropin releasing hormone (TRH) in normal

- man. J Clin Endocrinol Metab , 34 , 380-385.
363. Faglia, G. (1998). The clinical impact of the thyrotropin-releasing hormone test. Thyroid , 8 , 903-908.
364. Sawin, C.T., Hershman, J.M.& Chopra, I.J. (1977). The comparative effect of T4 and T3 on the TSH response to TRH in young adult men. J Clin Endocrinol Metab , 44 , 273-278.
365. Faglia, G., Beck-Peccoz, P., Ferrari, C., Ambrosi, B., Spada, A., Travaglini, P.& Paracchi, S. (1973). Plasma thyrotropin response to thyrotropin-releasing hormone in patients with pituitary and hypothalamic disorders. J Clin Endocrinol Metab , 37 , 595-601.
366. Mehta, A., Hindmarsh, P.C., Stanhope, R.G., Brain, C.E., Preece, M.A.& Dattani, M.T. (2003). Is the thyrotropin-releasing hormone test necessary in the diagnosis of central hypothyroidism in children. J Clin Endocrinol Metab , 88 , 5696-5703.
367. Borst, G.C., Osburne, R.C., O'Brian, J.T., Georges, L.P.& Burman, K.D. (1983). Fasting decreases thyrotropin responsiveness to thyrotropin-releasing hormone: a potential cause of misinterpretation of thyroid function tests in the critically ill. J Clin Endocrinol Metab , 57 , 380-383.
368. Gow, S.M., Elder, A., Caldwell, G., Bell, G., Seth, J., Sweeting, V.M., Toft, A.D.& Beckett, G.J. (1986). An improved approach to thyroid function testing in patients with non-thyroidal illness. Clin Chim Acta , 158 , 49-58.
369. Franklyn, J.A., Black, E.G., Betteridge, J.& Sheppard, M.C. (1994). Comparison of second and third generation methods for measurement of serum thyrotropin in patients with overt hyperthyroidism, patients receiving thyroxine therapy, and those with nonthyroidal illness. J Clin Endocrinol Metab , 78 , 1368-1371.
370. Toft, A.D., Irvine, W.J., Hunter, W.M., Ratcliffe, J.G.& Seth, J. (1974). Anomalous plasma TSH levels in patients developing hypothyroidism in the early months after 131I therapy for thyrotoxicosis. J Clin Endocrinol Metab , 39 , 607-609.
371. Toft, A.D., Irvine, W.J., McIntosh, D., Seth, J., Cameron, E.H.& Lidgard, G.P. (1976). Temporary hypothyroidism after surgical treatment of thyrotoxicosis. Lancet , 2 , 817-818.
372. Thein-Wai, W.& Larsen, P.R. (1980). Effects of weekly thyroxine administration on serum thyroxine, 3,5,3'-triiodothyronine, thyrotropin, and the thyrotropin response to thyrotropin-releasing hormone. J Clin Endocrinol Metab , 50 , 560-564.
373. Vagenakis, A.G., Braverman, L.E., Azizi, F., Portinay, G.I.& Ingbar, S.H. (1975). Recovery of pituitary thyrotropic function after withdrawal of prolonged thyroid-suppression therapy. N Engl J Med , 293 , 681-684.
374. Coya, R., Carro, E., Mallo, F.& Dieguez, C. (1997). Retinoic acid inhibits in vivo thyroid-stimulating hormone secretion. Life Sci , 60 , 247-250.
375. Sherman, S.I., Gopal, J., Haugen, B.R., Chiu, A.C., Whaley, K., Nowlakha, P.& Duvic, M. (1999). Central hypothyroidism associated with retinoid X receptor-selective ligands. N Engl J Med , 340 , 1075-1079.
376. Gow, S.M., Kellett, H.A., Seth, J., Sweeting, V.M., Toft, A.D.& Beckett, G.J. (1985). Limitations of new thyroid function tests in pregnancy. Clin Chim Acta , 152 , 325-333.
377. Emerson, C.H., Dysno, W.L.& Utiger, R.D. (1973). Serum thyrotropin and thyroxine concentrations in patients receiving lithium carbonate. J Clin Endocrinol Metab , 36 , 338-346.
378. Refetoff, S., Sunthornthepvarakul, T., Gottschalk, M.E.& Hayashi, Y. (1996). Resistance to

- thyrotropin and other abnormalities of the thyrotropin receptor. *Recent Prog Horm Res* , 51 , 97-120.
379. Abramowicz, M.J., Duprez, L., Parma, J., Vassart, G.& Heinrichs, C. (1997). Familial congenital hypothyroidism due to inactivating mutation of the thyrotropin receptor causing profound hypoplasia of the thyroid gland. *J Clin Invest* , 99 , 3018-3024.
380. Xie, J., Pannain, S., Pohlenz, J., Weiss, R.E., Moltz, K., Morlot, M., Asteria, C., Persani, L., Beck-Peccoz, P., Parma, J., Vassart, G.& Refetoff, S. (1997). Resistance to thyrotropin (TSH) in three families is not associated with mutations in the TSH receptor or TSH [see comments]. *J Clin Endocrinol Metab* , 82 , 3933-3940.
381. Ahlbom, B.D., Yaqoob, M., Larsson, A., Ilicki, A., Anneren, G.& Wadelius, C. (1997). Genetic and linkage analysis of familial congenital hypothyroidism: exclusion of linkage to the TSH receptor gene. *Hum Genet* , 99 , 186-190.
382. Boscato, L.M.& Stuart, M.C. (1988). Heterophilic antibodies: a problem for all immunoassays. *Clin Chem* , 34 , 27-33.
383. Despres, N.& Grant, A.M. (1998). Antibody interference in thyroid assays: a potential for clinical misinformation. *Clin Chem* , 44 , 440-454.
384. Silva, J.E.& Larsen, P.R. (1978). Peripheral metabolism of homologous thyrotropin in euthyroid and hypothyroid rats: acute effects of thyrotropin-releasing hormone, triiodothyronine, and thyroxine. *Endocrinology* , 102 , 1783-1796.